

# Mechanisms of Integral Membrane Protein Insertion and Folding

Florian Cymer<sup>1,†</sup>, Gunnar von Heijne<sup>1,2,†</sup> and Stephen H. White<sup>3,†</sup>

**1 - Center for Biomembrane Research, Department of Biochemistry and Biophysics, Stockholm University, SE-106 91 Stockholm, Sweden**

**2 - Science for Life Laboratory, Stockholm University, Box 1031, SE-171 21 Solna, Sweden**

**3 - Department of Physiology and Biophysics and the Center for Biomembrane Systems, University of California, Irvine, CA 92697, USA**

**Correspondence to Gunnar von Heijne:** Center for Biomembrane Research, Department of Biochemistry and Biophysics, Stockholm University, SE-106 91 Stockholm, Sweden. [gunnar@dbb.su.se](mailto:gunnar@dbb.su.se)

<http://dx.doi.org/10.1016/j.jmb.2014.09.014>

**Edited by F. Jacob-Dubuisson**

## Abstract

The biogenesis, folding, and structure of  $\alpha$ -helical membrane proteins (MPs) are important to understand because they underlie virtually all physiological processes in cells including key metabolic pathways, such as the respiratory chain and the photosystems, as well as the transport of solutes and signals across membranes. Nearly all MPs require translocons—often referred to as protein-conducting channels—for proper insertion into their target membrane. Remarkable progress toward understanding the structure and functioning of translocons has been made during the past decade. Here, we review and assess this progress critically. All available evidence indicates that MPs are equilibrium structures that achieve their final structural states by folding along thermodynamically controlled pathways. The main challenge for cells is the targeting and membrane insertion of highly hydrophobic amino acid sequences. Targeting and insertion are managed in cells principally by interactions between ribosomes and membrane-embedded translocons. Our review examines the biophysical and biological boundaries of MP insertion and the folding of polytopic MPs *in vivo*. A theme of the review is the under-appreciated role of basic thermodynamic principles in MP folding and assembly. Thermodynamics not only dictates the final folded structure but also is the driving force for the evolution of the ribosome–translocon system of assembly. We conclude the review with a perspective suggesting a new view of translocon-guided MP insertion.


© 2014 Elsevier Ltd. All rights reserved.

## Introduction

Membrane proteins (MPs) occur in all living cells and form part of key metabolic pathways, such as the respiratory chain and the photosystems. MPs also mediate the transport of solutes and the transduction of signals across membranes. For proper insertion into their target membrane, nearly all MPs require the aid of translocons, so-called protein-conducting channel proteins. In the past decade, structures of key translocon complexes that mediate the translocation or insertion of MPs have been solved and biochemical studies have yielded important insights into key aspects of MP insertion and folding. The cartoon in Fig. 1 illustrates a general scheme for the assembly of multi-span MPs. We will add many details to this scheme in the

course of this review, but many questions will still remain about the membrane insertion of proteins, how they fold into their final tertiary structure, and how they assemble into homo- and hetero-oligomeric complexes.

Our aim in this review is not to provide a comprehensive overview of the literature but rather to appraise critically current models of MP insertion and folding *in vivo* and to identify the most important gaps in our understanding of these processes. We start with a discussion of the basic biophysics of protein–lipid interactions and then turn our attention to how the biophysical principles are played out in the *in vivo* context. We suggest in the concluding section some outstanding questions that the field needs to address. We close with some general thoughts on how translocons may facilitate insertion



**Fig. 1.** This schematic cartoon represents in broad terms current thinking about the insertion of multi-span proteins into membranes. Two ideas are captured in the cartoon. First, TM segments (red) emerge from the ribosome and pass into the translocon (blue). Second, the nascent segments partition into the membrane from the translocon. As a starting point for discussion, we present alternative views of the MP insertion pathway in Fig. 15.

and folding. Through out this review, our focus is on helix-bundle integral MPs and SecYEG/Sec61-type translocons [found in the inner membrane of bacteria and the endoplasmic reticulum (ER) membrane of eukaryotic cells];  $\beta$ -barrel and peripheral MPs will not be discussed.


## Biophysical Boundaries of MP Insertion and Folding

### MPs and equilibrium


Molecular dynamics (MD) simulations of MPs in fluid lipid bilayers reveal graphically the complex environment of MPs. The environment of the SecYEG translocon is especially complex as a result of its large hour-glass-shaped interior filled with about 250 water molecules (Fig. 2). The structural stability of SecYEG and other MPs depends upon numerous physicochemical interactions, summarized in a simplistic way in Fig. 3. Despite the complexity, the equilibrium structures of MPs are in principle calculable given a complete quantitative description of the interactions. These interactions are understood in broad terms [1–3], but many essential details are lacking. Even though the physical chemistry of soluble proteins is also complex, we at least have a simple measure of their stability defined by their free energies of folding,  $\Delta G_{\text{fold}}$ . These free energies, usually obtained using calorimetric or chemical denaturation methods [4], are typically  $-5$  to  $-10$  kcal mol $^{-1}$  regardless of the number of amino acid residues [5].

Even defining what “unfolding” of an  $\alpha$ -helical MP means is problematic [6]. For example, bacteriorhodopsin (bR), a protein with seven transmembrane helices (TMHs), can be unfolded by heating, but unfolding is irreversible [7]. Other MPs can be unfolded reversibly with detergents, but the structure of the unfolded detergent–protein complex is unknown [8]. All we know for sure in either case is that the denatured proteins remain about 50% helical,

which means that  $\alpha$ -helices are extremely stable in lipids and detergents; this is expected from simple thermodynamic considerations (below). Despite the unsolved challenges of MP folding, understanding the thermodynamic stability of MPs is important because it defines the energetic framework within which the translocon apparatus must operate.



**Fig. 2.**  $\alpha$ -Helical MPs exist in their native state in highly thermally disordered lipid bilayers, as illustrated here for the SecYEG translocon [78] from *Methanococcus jannaschii*. The image is from an MD simulation of the translocon (PDB code 1RHZ) executed in a phospholipid bilayer. In this view, parallel with the membrane plane, the so-called gate helices 2b and 7 (red cylinders) were exposed by cutting away the lipid bilayer. Water molecules within the translocon are shown as van der Waals spheres in red (oxygen) and white (hydrogen). Waters surrounding the bilayer are shown as H-O-H bonds in blue-gray. Lipid acyl chains are white and the phospholipid headgroups are red. The image is provided courtesy of J. Alfredo Freites and Stephen H. White.



**Fig. 3.** Summary of the various interactions that stabilize MPs stably folded in fluid lipid bilayers (blue lines are interface boundaries, and red lines represent boundaries of the lipid hydrocarbon core). “Global bilayer effects” accounts for changes in the structure and stability of the lipid bilayer when perturbed by the protein [160], emphasizing that the bilayer itself sits in a free energy minimum determined by the tendency of the system to minimize exposure of the acyl chains (grey) to water (blue) on the one hand, as well as the tendency to maximize the distance between headgroups on the other. Both the bilayer and the protein must adjust structurally to minimize the free energy of the protein plus bilayer system. The protein shown (red helices) is bR determined to a resolution of 1.55 Å, PDB code 1C3W [161]. The image is modified from Ref. [162].

Thermodynamic measurements of MP stability are of little value unless native MPs are equilibrium structures. Several lines of evidence suggest that MPs are in fact equilibrium structures. Perhaps the single most important test of equilibrium is whether the MP structure is independent of the assembly pathway. Several experimental approaches, albeit for a limited number of MPs, are consistent with equilibrium. bR can be proteolytically cleaved into two membrane-spanning fragments and reconstituted separately into lipid vesicles. Fusion of the vesicles by multiple freeze–thaw cycles results in reconstitution of bR with a retinal absorption spectrum indistinguishable from wild-type spectrum [9]. Furthermore, low-resolution diffraction patterns of the reassembled protein are indistinguishable from those of wild-type protein [10]. If the final stages of folding MPs are guided by equilibrium thermodynamics, then co-expression of contiguous segments of an MP should lead *in vivo* to properly folded and functional proteins. This hypothesis has been confirmed for several  $\alpha$ -helical proteins, including rhodopsin [11], lactose permease [12], and the red cell anion exchanger protein [13]. Interestingly, even the  $\beta$ -barrel outer MP OmpA can be assembled from fragments [14]. Additional strong support for equilibrium comes from a recent study of the topology of lactose permease (LacY) in lipid bilayers [15],

which shows that, *in vivo*, LacY topology changes induced by manipulation of *Escherichia coli* inner membrane lipid composition can be replicated in lipid vesicles.


Given that MPs are equilibrium structures and ignoring possible kinetic barriers, we should, in principle, be able to reconstitute them into lipid vesicles or nanodiscs [16] without the intervention of a translocon under extremely dilute conditions: Imagine dropping a single MP into an aqueous phase containing lipid bilayer vesicles with an appropriate lipid composition. That single protein must eventually find its way to the vesicles and there fold spontaneously into the bilayer and eventually reach its native equilibrium state. This thought experiment is impractical to implement, of course, but an equivalent experiment would be to use a cell-free expression system containing only the essential components of protein synthesis. In essence, proteins emerging from the ribosome would find their way to lipid vesicles or nanodiscs and fold into the membrane in the absence of translocons. This approach seems to have been successful in several cases [17–21]. The main uncertainty is the exact composition of the cell-free extracts. It is not clear, for example, if the extracts are completely free of translocon components or chaperone components such as trigger factor [22].

### MP intrinsic interactions: TM helix folding and stability

In the absence of methods for folding and refolding  $\alpha$ -helical MPs reversibly in a structural context, the next best approach is to consider the interactions of fragments of MPs, such as  $\alpha$ -helices, with lipid bilayers. Because MPs are equilibrium structures, their intrinsic interactions with lipid bilayers can be described by any convenient set of experimentally accessible thermodynamic pathways, irrespective of the biological synthetic pathway. One particularly useful set of pathways is the so-called four-step model [1] (Fig. 4). Although these pathways do not necessarily mirror the actual biological assembly process of MPs, they can help us understand the thermodynamic constraints on MP structure formation and possibly give insights into the functioning of translocons.

Each of the steps in the four-step model has been intensively studied by several laboratories during the past 20 years [1–3,23]. Extensive measurements of water-to-bilayer and water-to-octanol partitioning free energies of model peptides [24–26] show that structure formation is dominated by the unfavorable free energy cost  $\Delta G_{\text{CONH}}$  of partitioning non-hydrogen-bonded peptide bonds and the favorable free energy of the hydrophobic effect  $\Delta G_{\text{HF}}$ . The latter free energy is readily calculated from the accessible surface area of non-polar amino acids, as shown long ago by Reynolds and Tanford [27]. They found that a





**Fig. 4.** A four-step thermodynamic cycle for describing the energetics of the partitioning, folding, insertion, and association of an  $\alpha$ -helix (red helices) in a lipid bilayer (grey). The process can follow an interfacial path, a water path, or a combination of the two. Studies of folding along the interfacial path are experimentally more tractable [163]. The  $\Delta G$  symbols indicate standard transfer free energies. The subscript terminology indicates a specific step in the cycle. The subscript letters are defined as follows: w, water; i, interface; h, hydrocarbon core; u, unfolded; f, folded; a, association. With these definitions, for example, the standard free energy of transfer from water to interface of an unfolded peptide would be  $\Delta G_{wiu}$ . However, some parts of the cycle (dashed box) are generally inaccessible experimentally. The image is modified from Ref. [1].


plot of transfer free energies against accessible surface areas  $A_{acc}$  of a large number of non-polar compounds yielded a linear curve with a slope of 20–25 cal mol<sup>-1</sup> Å<sup>-2</sup>, referred to as the atomic solvation parameter,  $\sigma$ , so that  $\Delta G_{HF} = \sigma A_{acc}$ . Measurements of the partitioning free energy of hydrophobic pentapeptides into *n*-octanol from water and careful accounting of their non-polar accessible surface areas yielded a value of -23 cal mol<sup>-1</sup> Å<sup>-2</sup> [24]. Interestingly, however, measurements of the partitioning free energies of the pentapeptides into the interface of phosphatidylcholine bilayers yielded a value of  $\sigma$  of about -11 cal mol<sup>-1</sup> Å<sup>-2</sup> [25], about one-half the octanol value. Why should the value be smaller? The hydrophobic effect depends upon complete dehydration of the non-polar surface when a solute is moved from bulk water to a bulk non-polar phase. The bilayer interface, however, contains waters with restricted motions due to headgroup hydration, as well as the interfacially exposed bilayer hydrocarbon core. A non-polar solute that partitions into the interface is never fully immersed in hydrocarbon and never fully dehydrated, causing  $\sigma$  to be reduced by half.

The favorable hydrophobic partitioning free energy of a non-polar amino acid is opposed by the unfavorable cost  $\Delta G_{CONH}$  of partitioning the peptide bond. However, if the peptide bonds can form hydrogen bonds (H-bonds) through secondary structure formation, the partitioning cost can be dramatically reduced [28,29]. The reduction occurs because  $\Delta G_{Hbond}$  of partitioning hydrogen-bonded peptide bonds is considerably lower than  $\Delta G_{CONH}$  of partitioning peptide bonds alone. For example, computational studies [30,31] of peptides in bulk alkanes suggest

that  $\Delta G_{CONH}$  for water-to-alkane transfer is +6.4 kcal mol<sup>-1</sup>, compared to only +2.1 kcal mol<sup>-1</sup> for  $\Delta G_{Hbond}$ . The per-residue free energy cost of disrupting H-bonds in an alkane is therefore about 4 kcal mol<sup>-1</sup>, which means that a 20-amino-acid transmembrane (TM) helix would cost in excess of 80 kcal mol<sup>-1</sup> to unfold within the membrane hydrocarbon. This explains why unfolded polypeptide chains cannot exist in a TM configuration.

The energetic cost of partitioning a peptide chain into the membrane interface is also reduced by H bonding to form secondary structure [28,29], the best current estimate being about -0.4 kcal mol<sup>-1</sup> per residue [32]. Although a modest number, the driving free energy for secondary structure formation by a 15-residue hydrophobic peptide would be about -6 kcal mol<sup>-1</sup> ( $\Delta G_{if}$  in Fig. 4). Because of this phenomenon, a modestly hydrophobic peptide such as 26-residue melittin from bee venom, which is largely unfolded in free solution, avidly folds into membrane interfaces [29].

The value of  $\Delta G_{Hbond}$  also sets the threshold for TMH stability. Even though hydrogen bonded, the cost of dehydrating peptide bonds upon insertion of a helix into the hydrocarbon core of a bilayer is still considerable, the best estimate being about +1.2 kcal mol<sup>-1</sup> per residue [33]. This explains why TMHs must be very hydrophobic: The unfavorable cost of partitioning the helical backbone must be compensated by the favorable cost of partitioning hydrophobic side chains. Figure 5 shows a free energy accounting for the insertion of the TM segment of glycophorin A (GpA), one of the first MPs to be sequenced [34]. The cost of partitioning the H-bonded helical backbone [ $\Delta G_{bb(h)}$ ] of



**Fig. 5.** The energetics of inserting an  $\alpha$ -helix into lipid bilayers (grey) is dominated by the peptide bonds, as illustrated here for the glycoporphin A (GpA) TM helix. Even with the helical backbone internally hydrogen bonded, it is costly to dehydrate the H-bonded peptide bonds ( $\Delta G_{bb(f)}$ ) upon insertion into the bilayer. For the helix to be stable, the favorable free energy of transfer of the side chains ( $\Delta G_{sc}$ ), determined by the hydrophobic effect, must compensate for the unfavorable  $\Delta G_{bb(f)}$ . In the case of GpA, the net stability of the helix  $\Delta G_{TM}$  is  $-12 \text{ kcal mol}^{-1}$ . The energetic cost of unfolding the polyglycine helix within the bilayer is immense:  $\Delta G_{bb(u)}$  is greater than  $100 \text{ kcal mol}^{-1}$ ! Modified from Ref. [162].

$+24 \text{ kcal mol}^{-1}$  is well compensated by the favorable hydrophobic free energy of  $-36 \text{ kcal mol}^{-1}$  for partitioning the side chains ( $\Delta G_{sc}$ ). The net free energy  $\Delta G_{TM}$  favoring insertion is thus  $-12 \text{ kcal mol}^{-1}$ . As is common in so many biological equilibria, the free energy minimum is the small difference of two relatively large opposing energetic terms. This simple example emphasizes that the balance between the unfavorable cost of dehydrating the H-bonded peptide bond and the favorable gain from dehydrating non-polar side chains determines whether or not an  $\alpha$ -helix is independently stable in a TM configuration. The cost of partitioning the unfolded GpA peptide backbone would exceed  $100 \text{ kcal mol}^{-1}$  to emphasize again the importance of H-bonds in determining structure! This huge cost explains why heat or chemical denaturation of MPs always results in “unfolded” states that are rich in  $\alpha$ -helical structure.

But is the cost realistic? Atomic force microscopy (AFM) provides an answer. TMHs can be pulled out of membranes one by one using AFM methods [35]. In a classic AFM study of bR [36], for example, a long, flexible cantilever with an atomically sharp probe attached to one end was used to scan the surface of a native bR membrane. The deflection of the cantilever as it moved across the surface was recorded, producing a topographic map of the surface. The AFM was then used to pull single molecules out of the surface while recording the force exerted on the cantilever. Because both the

force  $F$  on the cantilever and the distance  $d$  between the tip and the membrane can be measured simultaneously, the unfolding of the protein can be visualized as a force–distance curve. Importantly, the unfolded state is an extended polypeptide chain in aqueous buffer. This means that the underlying thermodynamic states are the folded protein in the bilayer and the unfolded protein in water. The unfolding is a non-equilibrium process and is not reversible, but the data are nevertheless useful because the non-equilibrium problem can be worked around using Jarzynski's identity [37]. Careful application of the identity to force–distance curves collected over wide range of temperatures and pulling rates yielded [38], for the first five helices of bR, a free energy difference between membrane-folded and water-unfolded states of  $230 \pm 40 \text{ kcal mol}^{-1}$  or an average free energy change of about  $1.3 \text{ kcal mol}^{-1}$  per residue. This value agrees well with the value of about  $200 \text{ kcal mol}^{-1}$  computed from octanol–water partitioning free energies [1,24], which underlie the calculations of Fig. 5.

Apart from these AFM measurements that apply in special cases, essential thermodynamic information is generally difficult to obtain through direct measurements because greasy peptides are very insoluble and have a strong tendency to aggregate in water. An alternative for understanding in atomic detail the partitioning of highly hydrophobic peptides is to use microsecond-scale equilibrium MD simulations [39].

Simulation technology has not yet advanced sufficiently to examine the folding of multi-span MPs, but the technology is good enough to examine the folding of single-span polyleucine segments. Such simulations are initiated by placing a single unfolded polyleucine peptide into a simulation cell containing a lipid bilayer and allowing the simulation to run until all accessible states have been sampled many times and cataloged. In simulations lasting several microseconds, polyleucine peptides are observed to adsorb into the membrane interface where they fold into a helical conformation and make numerous excursions into the membrane as a TM helix. Figure 6 shows a representation of the folding and insertion of a Leu10 peptide in which the insertion depth and helicity of the peptide are sampled during the course of the simulation. For each sampled time point, the insertion depth (relative the bilayer center) is plotted against the helicity, and the points are connected sequentially by lines.

Two observations are significant. First, the peptide rapidly adsorbs at the bilayer interface. Second, after the first 50 ns, the peptide becomes  $\alpha$ -helical and moves repeatedly between a helical surface configuration and a helical TM configuration. This behavior is exactly what one would expect from considerations of the cost of peptide bond partitioning described above (Figs. 4 and 5). During the time course of the simulation, the peptide is never seen to leave the membrane and return to the aqueous phase, which is the behavior expected from extensive studies of the interactions of melittin with bilayers (above). Indeed, microsecond-scale simulations of melittin folding at the membrane interface are in quantitative agreement with experiments [40,41].


These considerations of the partitioning of peptides into membranes provide an important perspective on

how translocons may manage the MP assembly reaction pathway. They lead inexorably to the conclusion that the membrane interface in the vicinity of the translocon likely participates in assembly.

### MP intrinsic interactions: Helix–helix interactions


There is much more to intrinsic interactions than just the folding and insertion of single TMHs into lipid bilayers because so many biologically important MPs are multi-spanning proteins. The other part of the MP folding problem is therefore the energetics of helix–helix interactions that cause single helices to come together to form functional three-dimensional structures in the lipid bilayer. Popot and Engelman first focused attention on this problem through their “2-stage model” of folding [42], which was an outgrowth of their work on the reassembly of functional bR from proteolytically cleaved fragments separately reconstituted into lipid vesicles. Our clearest understanding of helix–helix interactions comes from studies of glycophorin A (GpA) that forms dimers when solubilized in sodium dodecyl sulfate (SDS) [43] and lipid bilayers [44]. Engelman *et al.* took advantage of this observation to discover the location and properties of the dimerization domain,  $-L_{75}IxxG_{79}V_{80}xxG_{83}V_{84}xxT_{87}-$ , within the GpA TMH [45,46] and to determine the structure of the dimer in detergent micelles using NMR [47] (Fig. 7). Langosch *et al.* extended GpA dimerization measurements to *E. coli* inner membranes [48].

The GpA dimer is held together by van der Waals interactions and tight shape complementarity, called “knobs-into-holes packing” [49,50]. The GpA data led to the identification of the first



**Fig. 6.** Equilibrium microsecond-scale simulations of the folding and membrane insertion of polyleucine sequences (here 10 leucines) reveal only two states [39]. The simulation shown here begins with the unfolded peptide (U) in water (W) about 10 Å from the phosphatidylcholine bilayer (grey). Within a few nanoseconds (ns), it adsorbs to the membrane interface and never returns to the bulk water phase. After the next 40 ns, the peptide becomes  $\alpha$ -helical and fluctuates between being on the surface (S) and across the membrane (TM) during the ensuing several microseconds. The trajectory of the simulation is represented as a plot of the insertion depth of the

peptide's center of mass against its helicity. The sampled time points are connected sequentially by blue lines. Modified from Ref. [39]. This simulation reveals the importance of the membrane interface in MP folding and suggests that the interface may play a role in translocon-guided insertion of TM helices.



**Fig. 7.** Helix–helix interactions of the glycophorin A (GpA) dimer (blue and orange helices) based upon the structure of GpA in SDS [47]. (a) Glycine (or other small residues) separated by three residues allows the helices to pack tightly. (b) Other amino acids in the structural vicinity of the GXXXG motif can facilitate or inhibit specific binding of complementary surfaces [120].

widespread interaction motif for TMHs, the so-called GXXXG motif [51,52]. The interaction free energies of GpA dimer formation are readily determined for detergent-solubilized helices using analytical ultracentrifugation to measure the equilibrium amounts of monomers and dimers [53], but the relationship between interaction free energies in detergents and in lipid bilayers has been unclear. A recent study shows that helix dimer stability in lipid bilayers depends strongly on the type of lipid, ranging from  $-12 \text{ kcal mol}^{-1}$  in pure phosphatidylcholine bilayers to about  $-3 \text{ kcal mol}^{-1}$  in *E. coli* lipids [54], the latter value being similar to that observed in natural membrane “blebs” [55].

H-bonds between TMHs can also be important for stabilization. For example, TM segments composed only of leucine have relatively little tendency to dimerize *in vivo* in *E. coli* unless a polar residue, especially Asn, Asp, or Glu, is also present in the sequence to form H-bonds [56]. However, a double-mutant thermodynamic cycle analysis of bR has shown that the stabilization free energy of single side-chain H-bonds contributes only about  $0.6 \text{ kcal mol}^{-1}$ . Of course, if multiple H-bonds are present between side chains in an MP, the additive stabilization could become significant.

Because of the simple thermodynamic considerations discussed earlier, MPs are absolutely constrained to have their native helical structure in bilayer membranes. The first stage of the 2-stage model [42]—the establishment of TMHs across membranes—is accomplished through co-translational insertion and the catalytic activity of the translocon (below). Cells thus avoid the thermodynamic challenges associated with managing highly non-polar proteins in the aqueous environment. The second stage of folding—helix association

within the bilayer environment—should be the primary focus of MP stability rather than complete denaturation of MPs using heat or detergents, which are problematic. Just as we measure the stability of soluble proteins in their native aqueous environment, we should really measure the stability of MPs within their native membrane environment. The question is then simply the energetic cost of separating the TMHs from one another within a lipid bilayer. This challenging task has now been accomplished for bR in a bicelle environment using so-called steric trapping [57]. Steric trapping is based upon engineering two streptavidin binding sites into the protein that are close together in the native state. For folded proteins, only a single monovalent streptavidin can bind due to steric overlap of the binding sites. However, when the protein is unfolded in the membrane, then streptavidin can bind to both sites. The free energy of unfolding can be determined by measuring of the amount of doubly bound protein during unfolding. In this way, Chang *et al.* determined a value of approximately  $-11 \text{ kcal mol}^{-1}$  for the unfolding of bR [58], which is within range of values observed for water-soluble proteins. This is not a marginal value of stability. If other MPs are eventually found to have values similar to those of water-soluble proteins and membrane-bound bR, then one can safely assume that native three-dimensional structure will follow once TMHs are inserted into membranes.

## Biological Boundaries of MP Biogenesis

How does the cell exploit and control the physical interactions that underlie the spontaneous insertion of hydrophobic peptides into lipid bilayers in order to produce properly folded helix-bundle MPs while avoiding the ever-present risk of protein aggregation? The basic solution to this problem that nature has come up with is co-translational insertion: With few exceptions [59], helix-bundle MPs are inserted directly into a target membrane as they come out of the ribosome. Co-translational insertion is mediated by one or other type of translocon, that is, a protein conduction channel in the membrane that is constructed such that it can both shield polar parts of the polypeptide from lipid contact as they are translocated across the membrane and expose at the same time hydrophobic segments in the protein to the surrounding lipid bilayer, facilitating the formation of TMHs.

In this section, we first review recent advances in our understanding of how ribosomes from different kinds of organisms handle nascent polypeptide chains and how ribosomes synthesizing MPs are specifically targeted to translocons. We then discuss current models of translocon-mediated insertion and folding of MPs.



## Ribosomes

Ribosomes are highly conserved RNA–protein complexes that translate mRNA into protein in all living cells. For our present purposes, the most important parts of the ribosome are the polypeptide transferase center (PTC) where the incoming tRNA delivers its amino acid to the growing polypeptide, the tunnel in the large ribosomal subunit through which the nascent chain exits the ribosome, and the polypeptide exit site on the “backside” of the large subunit (Fig. 8). Peptide bond formation at the PTC is catalyzed by ribosomal RNA (“The ribosome is a ribozyme” [60,61]). Large sections of the tunnel walls are also formed by ribosomal RNA, creating a polar, negatively charged environment for the nascent chain. There is a higher density of ribosomal proteins around the exit site, and many accessory proteins such as chaperones and targeting factors bind there. The area around the exit site is also involved in docking the ribosome to the translocon, such that the nascent chain can pass directly from the ribosomal tunnel into the translocon channel.


The chief differences between prokaryotic and eukaryotic ribosomes are found in their protein composition, with eukaryotic ribosomes having a larger complement of proteins than prokaryotic ones. There are also some extra “expansion loops” in the RNA present in eukaryotic ribosomes [62–64]. The core ribosomal structure is highly conserved, how-

ever, and the basic operational principles—including the biosynthesis of integral MPs—are very similar from bacteria to man.

The key player in the early steps of MP biosynthesis is the signal recognition particle (SRP). In eubacteria such as *E. coli*, SRP is composed of a small 4.5S RNA and a single protein subunit, Ffh [65,66]. Ffh contains both a signal-peptide binding domain and a GTPase domain, and it can bind near the polypeptide exit site on the ribosome. When a sufficiently hydrophobic polypeptide segment, typically a signal peptide or a segment that will eventually form a TMH, appears at the exit site of the ribosome-nascent chain complex, it binds to SRP with subnanomolar affinity [67]. Binding will cause a major structural rearrangement of the SRP [66,68], priming the ribosome-nascent chain–SRP complex for binding to the SRP receptor FtsY located on the surface of the inner membrane. The ribosome–SRP–FtsY complex then binds to the SecYEG translocon (see the next section). FtsY also has a GTPase domain, and proper docking of SRP to FtsY brings the SRP and FtsY GTPase domains into juxtaposition such that the two complete active sites are formed and two GTPs can be hydrolyzed. This process, which serves as a quality control mechanism by assuring that the ribosome is correctly docked to SecYEG [69], leads to the release of the nascent chain from SRP, allowing it to insert into the translocon.

The mammalian SRP is more complex, with a larger 7S RNA and five additional proteins. The Ffh homologue is called SRP54 and has a similar signal-peptide binding groove [65,66,70]. A major difference between the bacterial SRP and the mammalian SRP is that the latter not only binds hydrophobic segments in the nascent chain and mediates targeting to the translocon but also slows down or halts translation when bound to a ribosome-nascent chain complex [65]. Presumably, this gives the ribosome more time to find and dock to a translocon in the complex cytosolic environment of a eukaryotic cell, ensuring the tight coupling between translation and membrane insertion.

Residues in the nascent polypeptide chain can interact with the tunnel wall, especially in its upper parts. Such interactions can induce translational stalling, presumably by subtly altering the geometry of the PTC. Specific peptide sequences, called arrest peptides (AP), can stall translation on their own (intrinsic APs) or in combination with a small molecule such as an amino acid or an antibiotic (inducible APs) [71]. The latter are important in regulation of antibiotic resistance genes, which have APs that are activated in the presence of the antibiotic [72]. As will be explained below, translational APs have recently been employed as “force sensors” to detect dynamic interactions between a growing nascent chain and its immediate surroundings.



**Fig. 8.** High-resolution structure of a mammalian ribosome–Sec61 complex. The structure was obtained using advanced cryo-EM methods. PTC indicates the peptidyl transferase center. The image is from Voorhees *et al.* [64].



### Translocon-mediated membrane insertion

In bacteria, membrane insertion of nearly all helix-bundle MPs is mediated by two translocons: SecYEG and the structurally unrelated YidC. Some proteins require only one or the other, and some require both. YidC is about 5-fold more abundant in *E. coli* than SecYEG [73,74]. SecYEG and YidC can form a 1:1:1:1 super-complex (the holo-translocon) together with SecDF (4-fold lower cellular abundance than SecYEG) and YajC (3-fold higher cellular abundance than SecYEG) [75].

The eukaryotic Sec61 translocon in the ER is homologous to SecYEG, but there are no YidC, SecDF, or YajC homologues in the ER. YidC homologues are, however, found in the inner mitochondrial membrane (Oxa1) and the chloroplast thylakoid membrane (Alb3/Alb4) [74]. On the other hand, the Sec61 super-complex involves additional MPs such as TRAM, TRAP, and the oligosaccharyl transferase that have no counterparts in *E. coli*. Eukaryotic Sec complexes also include Sec62 and Sec63, that facilitate post-translational transport (reviewed in Ref. [75]). Mitochondria in most organisms do not have an equivalent to Sec61 in the inner membrane; the only translocase there is Oxa1. Because of the absence of a Sec-type translocon, mitochondrial ribosomes are specialized to bind directly to the inner membrane [76].


The SecYEG/Sec61 translocon can mediate both co- and post-translational translocations of proteins; in bacteria, most secreted proteins are translocated post-translationally with the help of the SecA motor protein, whereas MPs are nearly always inserted into the membrane co-translationally [77]. In contrast,

YidC mediates post-translational membrane insertion, except in mitochondria where a C-terminal domain in Oxa1 can bind to mitochondrial ribosomes in a co-translational fashion [74]. In eukaryotic cells, insertion of proteins into the ER membrane is nearly always co-translational, while translocation of secretory proteins into the lumen of the ER can be both co-translational and post-translational [77].

High-resolution three-dimensional structures (Fig. 9) are known for several SecYEG translocons from different prokaryotic organisms [78–80] and for the Sec61 complex from pig [64]. The central translocation channel is formed by the SecY subunit (Sec61 $\alpha$  in eukaryotes), a protein with 10 TMHs. A small plug domain closes the channel from the periplasmic side, presumably in order to prevent ion leakage through the inactive translocon. The SecY subunit is buttressed by SecE that partly encircles SecY and enhances its stability [81]. SecG is more peripherally located, and its precise function is unclear [82].

A lateral gate located between TMH2b and TMH7 in SecY can open up the channel toward the surrounding membrane, as seen in Fig. 9. The simplest conceptual model for how TMHs in the nascent polypeptide are inserted into the membrane is that they first enter the central SecYEG channel, then exit it via the lateral gate and insert into the lipid bilayer one by one (Fig. 1). In essence, they partition between the translocon and the membrane. As will be shown below, this is certainly an oversimplification, but it serves as a good starting point for the discussion.

The idea of translocon/membrane partitioning can be examined experimentally by suitably designed test proteins in which a hydrophobic segment is placed far



**Fig. 9.** Structure of the SecYE translocon from *Pyrococcus furiosus* [80]. The left panel shows a view in the plane of the membrane, and the right panel shows a view from the cytoplasm. The plug domain is circled, and the residues in the hydrophobic ring are shown as van der Waals spheres, indicated by asterisks (\*). The arrow points into the lateral gate between TMH2b and TMH7. SecE is shown in red, and SecY is color coded from N-terminus (blue) to C-terminus (orange).


away from any potentially interfering TMHs. Assays to measure the efficiency of membrane insertion of such model hydrophobic segments (H-segments) have been developed, both for Sec61-mediated insertion into the mammalian ER [83,84] and SecYEG-mediated insertion into the inner membrane of *E. coli* [85]. Analysis of hundreds of H-segments with systematically varied amino acid sequences led to the establishment of a “biological” hydrophobicity scale, in which the contribution from each of the 20 natural amino acids to the overall apparent free energy of membrane insertion of the H-segment ( $\Delta G_{\text{app}}$ ) is given as a function of the position of the residue in the H-segment [83]. Further studies in which an amino acid with a non-natural side chain—either a linear alkyl chain with 1 to 8 carbons or aromatic groups as large as biphenyl—was included in different position in the H-segment made it possible to measure the contribution of apolar surface area (i.e., hydrophobic effect with solvation parameter  $\sigma$ ) to  $\Delta G_{\text{app}}$ : For an aliphatic surface,  $\sigma$  was determined to be  $-10 \text{ cal mol}^{-1} \text{ \AA}^{-2}$ , and for aromatic surface, the value was  $-7 \text{ cal mol}^{-1} \text{ \AA}^{-2}$  [86] when at the center of the H-segment. These solvation parameter values, measured for the translocon-mediated insertion process, are a factor 2 to 3 smaller than expected for simple water-to-hydrocarbon transfer free energies. Intriguingly, however, these values are similar to solvation parameters determined for the partitioning of hydrophobic peptides into bilayer interfaces [25] (see **MP intrinsic interactions: TM helix folding and stability**, above). The similarity suggests that the state of water in the translocon may be similar to the state in hydrated bilayer headgroups [87]. The smaller solvation parameter obtained in the translocon partitioning experiments [86] thus raises the possibility that the nascent chain may be exposed to the surface of the lipid bilayer surrounding the translocon at an early stage of membrane insertion. This possibility has recently received support from equilibrium MD simulations of the water-to-bilayer partitioning of H-segments whose values of  $\Delta G_{\text{app}}$  were determined experimentally [88].

Recently, a structure of a ribosome-nascent chain-translocon complex has been obtained at 3.4 Å resolution [64] (Fig. 8); and two more, at 8–15 Å resolution [89,90], all by single-particle cryo-electron microscopy (EM). The interpretation of the structures is complicated, however, because they represent equilibrium configurations—established using various experimental protocols—that may or may not represent way points on the physiological insertion pathway. A particular problem is that the translocons in the structures are invariably in detergent micelles rather than lipid bilayers.

The first structure [64] provides detailed views of both an idle ribosome-translocon complex with no nascent chain and an active complex in which the nascent chain can be followed through the ribosome but is not visible within the translocon. The cytoplasmic

end of the lateral gate is partly open in the active complex, and the plug domain is displaced from its original location. In the second structure [89], an H-segment engineered into a host protein has been caught in an open lateral gate, partly exposed to the surrounding detergent, and in the third [90], the hydrophobic signal peptide of the periplasmic protein DsbA is similarly located. These structures show that a TMH can occupy the lateral gate, but these do not address the question of whether such H-segments first enter the central channel in the translocon and then exit into the lipid bilayer via the lateral gate, slide progressively into the lateral gate from the top, always being partly exposed to lipid (Fig. 10), or perhaps first come into direct contact with the lipid bilayer. If the energetic barrier to opening the lateral gate is not too high, the “sliding” possibility seems more likely to represent the lowest-energy trajectory. Indeed, this is how nascent chains behave in a coarse-grained MD model of translocon-mediated protein translocation and insertion developed by Zhang and Miller [91]. A recent crosslinking study has identified a possible initial binding site for TMHs at the top of the lateral gate and in close proximity to surrounding lipid [92], from which the TMH could conceivably slide deeper into the membrane within a progressively expanding lateral gate. In the sliding model, a TMH would be partially exposed to lipid already at an early stage during insertion, possibly explaining the coincidence between insertion free energies calculated from the biological hydrophobicity scale and those obtained by MD for peptide insertion into a bilayer alluded to above. Further, the sliding model would predict that highly hydrophobic TMHs may insert directly into the membrane while making little contact with the lateral gate, amphiphilic TMHs insert with their more polar face buried in the lateral gate, and marginally hydrophobic TMHs—i.e., TMHs that insert efficiently only by making specific interactions with already inserted TMHs [93–95]—would insert largely through the translocon channel and only expose a limited surface through the lateral gate.


A high-resolution structure of the YidC translocon has just been published [96], as has an accurate structural model [97]. Somewhat surprisingly, YidC is a monomer in the crystal, while a previous projection structure obtained by electron crystallography of two-dimensional crystals showed it as a dimer [98], and the density attributed to YidC in a single-particle EM study of ribosome-YidC complexes was also interpreted as representing a YidC dimer [99]. On the other hand, and consistent with the X-ray structure, binding studies using YidC reconstituted into nanodiscs suggested that ribosome-nascent chain complexes bind only a single copy of YidC [100]. Very recent cryo-EM structures of YidC bound to a ribosome are also consistent with a monomer [97,101].



**Fig. 10.** Two models for translocon-mediated insertion of a  $N_{\text{out}}-C_{\text{in}}$  orientated TMH in a single-span (type I) MP. (a) The "in-out" model. The TMH (in black) first moves all the way into the central translocon channel and then exits sideways through the lateral gate. (b) The "sliding" model. The TMH slides along the lateral gate into the membrane, with one side exposed to lipid at all times. The leading polar segment penetrates through the lateral gate and is shielded from lipid contact. SecE is shown in red, and SecY is color coded from N-terminus (blue) to C-terminus (orange). On the right-hand side, a schematic of the translocon (blue) interacting with an MP (green) and a membrane inserting TMH (red) are shown.

The X-ray structure and accompanying MD simulations show a deep, hydrated cleft within YidC that extends halfway across the membrane and is open toward both the cytoplasm and the lipid bilayer (Fig. 11a). The cleft is capped on the periplasmic side by what appears to be a tightly

packed, stable structural domain. When Trp244 at the top of the cleft is substituted by the photo-activatable amino acid analog *p*-benzoyl-L-phenylalanine, it can be crosslinked *in vivo* to a co-expressed substrate protein, showing that substrate has access to the deep cleft [96]. There is also a



**Fig. 11.** Structure of the YidC translocon from *Bacillus halodurans* [96] viewed in the plane of the membrane, looking into the hydrated cleft in the center of the protein. The left panel shows a surface representation; the right panel shows the walls of the cavity, with the rest of the protein in stick representation.



rather mobile part, composed of two  $\alpha$ -helices that lie flat on the cytoplasmic face of the membrane, in front of the entrance to the cleft. The role of this part is unclear; perhaps it facilitates membrane insertion of substrate TMHs by locally distorting the lipid bilayer.

The structure [97,101] is suggestive of an insertion mechanism in which a short polar tail or loop in the substrate protein penetrates at least halfway across the membrane within the water-filled cleft, leaving the adjoining hydrophobic TMHs in the lipid bilayer, similar to the sliding model proposed for SecYEG above. The resulting intermediate state would be of lower free energy than an initial state where the TMH is embedded in the lipid headgroup region, but it would be of higher free energy than a final state where the TMH completely spans the membrane. According to this model, YidC is designed to lower the energy barrier for the initial, partial insertion of the TMH. A speculative possibility is that the outer lipid monolayer becomes sufficiently perturbed in the vicinity of the intermediate YidC–substrate complex to make it possible for the polar tail or loop to translocate fully across the membrane at a reasonable rate.


A comparative analysis of the insertion of model TMHs into the inner membrane of *E. coli* of YidC- and SecYEG-dependent MPs showed that the threshold hydrophobicity required for 50% insertion of a hydrophobic segment of composition  $nL/(19-n)A$  is similar ( $n_{50\%} \approx 1-2$ ) for the two translocons [85,102]. Likewise, the individual contributions to the overall  $\Delta G_{app}$  for YidC- and SecYEG-dependent insertions are similar for the non-polar residues, whereas polar and charged residues have roughly 2-fold larger  $\Delta G_{app}$  values when membrane insertion is mediated by the YidC translocon (Fig. 12). Possibly, a TMH is more lipid exposed during YidC-mediated

insertion than during insertion mediated by the SecYEG translocon (where polar residues may be sequestered into the lateral gate region, away from lipid contact).

Because some fraction of SecYEG and YidC are bound together in the holo-translocon, they can conceivably act sequentially on MP substrates or may even act simultaneously on different parts of a multi-spanning MP. Within the context of the holo-translocon, where the lateral gates in SecYEG and YidC conceivably are not far apart, one can imagine that an incoming N-terminal TMH, or an internal “helical hairpin” composed of two TMHs with a short connection loop, can preferentially partition into the membrane via YidC, while TMHs flanked by longer polar segments that cannot fit into the YidC cavity will preferentially be inserted via SecYEG.

### Ribosome–translocon–membrane insertion pathways

The progression of a hydrophobic TM segment along the ribosome–translocon–membrane insertion pathway has been followed mainly by chemical crosslinking experiments and, more recently, by using translational APs as *in vivo* force sensors. Many crosslinking experiments have been carried out by *in vitro* generation of stalled ribosome–nascent chain–translocon complexes using dog pancreas rough microsomes, that is, ER-derived membrane vesicles that contain Sec61 translocons. Early studies showed that signal peptides in stalled nascent chains could be crosslinked both to translocon components and to lipid [103], suggesting that the signal peptide is transiently held in an interfacial location between the translocon and surrounding lipid. This is in full agreement with the EM structures discussed above.



**Fig. 12.** Amino acid  $\Delta G_{app}$  values for Sec61-, SecYEG-, and YidC-dependent membrane insertions of a model TMH [84,85,102]. The SecYEG (a) and YidC (b) data are plotted against the Sec61 data. Full lines indicate linear fits to the data (with the equations given in the panels), while the broken line in (b) is the linear fit obtained when including only non-polar and weakly polar residues (slope = 0.8) [85].


As noted above, a recent crosslinking study has further identified an early interaction between N-terminal TMHs in a substrate protein with residues located at the cytoplasmic tip of the lateral gate in Sec61 [92], as would be expected if hydrophobic segments start to partition into the lateral gate region immediately upon entering the translocon rather than first moving into the central channel and exiting through the lateral gate only at a later stage (cf., Fig. 10b).

The development of a novel technique where translational APs are used to measure forces acting on a nascent chain during co-translational processes such as membrane insertion now makes it possible to study the kinetics of MP insertion and folding *in vivo*. APs are short stretches of polypeptide, typically ~10–15 residues long, that bind in the upper parts of the ribosomal tunnel and induce ribosomal stalling at a specific codon in the mRNA [71]. Stalling can be prevented if a sufficiently strong “pulling force” acts on the nascent chain at the precise point when the ribosome reaches the critical codon [104]; presum-

ably, pulling on the nascent chain breaks the interactions between the AP and the ribosome that control stalling.

APs from the SecM protein have proven particularly useful. The AP from *E. coli* SecM is 17 residues long and rather weak, but stronger APs have been found in SecM proteins from other bacteria [105]. The strongest AP known to date is a mutated version of the SecM AP from *Mannheimia succiniciproducens* with sequence HPPIRGSP (called Ms-Sup1) [106]. This short AP can be introduced into any protein and will report on the tension in the nascent chain at the precise point during translation when the ribosome reaches the proline codon at the 3' end of the AP.

The SecYEG-mediated insertion of polypeptide segments (H-segments) of varying hydrophobicity into the inner membrane of live *E. coli* cells was recently analyzed using the approach detailed in Fig. 13a [106]. Strong pulling forces proportional to the hydrophobicity of the H-segment were recorded at



**Fig. 13.** Using arrest peptides (APs) as *in vivo* force sensors [106]. (a) An AP (AP; in blue) is inserted into an MP with two natural TMHs (TM1, TM2; in black) and a model TMH composed of six leucines and 13 alanines (H; in red). Depending on the length  $L$  of the tether between the H-segment and the AP, the H-segment will be in different locations relative to the translocon at the time when the ribosome reaches the last codon in the AP, as shown in the cartoon. The pulling force  $F(L)$  on the nascent chain will determine the fraction of stalled *versus* full-length protein produced. The fraction of full-length protein can be determined by [ $^{35}$ S]Met pulse-labeling of growing *E. coli*, followed by immunoprecipitation of the protein construct and analysis by SDS-PAGE, as shown on the right for a construct with  $L = 63$  residues (a control construct in which a critical proline in the AP has been mutated to alanine, preventing stalling, is also shown). (b) Fraction full-length protein as a function of  $L$  for a set of constructs designed as shown in (a) (top).

tether lengths  $L \approx 30$  and  $L \approx 40$  residues (Fig. 13b). From the known dimensions of the ribosome–translocon complex, the peak in the force profile at  $L \approx 40$  residues most likely corresponds to the final insertion of the H-segment into the membrane. The interaction responsible for the peak at  $L \approx 30$  residues is more difficult to pinpoint: It depends only on the hydrophobicity of the N-terminal end of the H-segment and could represent an interaction between the H-segment and the cytoplasmic face of the lipid bilayer or possibly an interaction akin to the one identified between an N-terminal TMH and the tip of the lateral gate in Sec61 discussed above [92]. Regardless, the data show that APs can be used to measure forces acting co-translationally on a nascent chain with high sensitivity and high spatial precision (a force profile recorded with single-residue precision such as the one shown in Fig. 13b has a spatial resolution corresponding to one residue in an extended conformation, i.e.,  $\sim 3$  Å).

Segments that flank the TMHs can also affect membrane insertion. In particular, positively charged residues at the ends of a hydrophobic segment can either increase or decrease its insertion propensity, in accordance with the “positive-inside” rule [107–110].

### Folding of Polytopic MPs *In Vivo*

Translocon complexes mediate the insertion of TM segments into membranes. Although this process is guided by the amino acid composition and structure of the segment, its interaction with the translocon and translocon accessory components can be critical. As the polypeptide chain moves through the translocon, it is continuously scanned. A ring of hydrophobic residues is present in the middle of the channel in both SecYEG and Sec61 translocons [78,111] (Fig. 10). The interaction of the nascent chain with this ring of hydrophobic residues might be involved in the opening of the lateral gate and the transfer of TM segments between the translocon and membrane [112,113], where they can be exposed to the lipid environment. If a segment is hydrophobic enough, it will insert into the membrane, and current models assume that the TM segments in polytopic MPs are inserted sequentially [114]. One would expect that replacement of the hydrophobic residues in the ring would destabilize the translocon, but that is not the case. Replacement with polar, even charged, residues were not destabilizing [111]. The replacement does, however, affect  $\Delta G_{\text{app}}$  for membrane insertion of H-segments, consistent with the ring playing a role in translocon/membrane partitioning [112].

As discussed earlier in MP intrinsic interactions, the  $\alpha$ -helical structure of a nascent TMH forms at the latest during the exposure of the segment to the lipid bilayer, driven by the necessity to shield the polar backbone from an unfavorable exposure to the

hydrocarbon core of the membrane. In case of interactions with the translocon channel, a helical structure apparently can be induced even before the exposure to a lipid bilayer [92]. Helical structure may even be formed in the ribosome exit tunnel [115–118]. The stage at which an  $\alpha$ -helical structure develops can be important for the formation of tertiary structure, because amino acid side chains are positioned on different faces of an  $\alpha$ -helix, thereby allowing favorable alignment with residues in other TM segments to form tertiary structure.

In polytopic MPs, individual TMHs can form interactions that determine the final fold and function of the mature protein. As discussed earlier, the interactions between TM segments can consist of distinct interaction motifs, such as the GxxxG motif of GpA discussed above (Fig. 7). More generally, small residues spaced four residues apart (small-XXX-small) represent a prominent motif that allows TMHs to approach each other closely and pack tightly due to van der Waals attraction and steric constraints. Extensions of this motif (small-XXX-small-XXX-small) allows for flexibilities of two helices as multiple small residues, each spaced in the distance of four, can form a groove in the surface of  $\alpha$ -helices [119]. While any two helices containing such a motif could approach each other in principle, residues surrounding these motifs can provide certain specificity to a given interaction [120–122].

Although these motifs are highly abundant in MPs [123], many other types of interactions that require more specificity can be formed between TMHs, interactions with multiple TMHs at the same time or a high degree of flexibility. However, many interactions between TMHs cannot be explained by these simple interaction motifs [124]. Highly specific interaction networks between multiple TMHs that determine the final structure can be formed, and the flexibility between these interactions allows for tertiary structure changes that are critical for function. It is therefore not surprising that simple interaction motifs occur mainly in rather simple function contexts such as the on/off function of some receptors [125].

Many MPs have more complex TM domains, for example, TM channels. Here, the TMHs must form an aqueous pore, lined by hydrophilic residues within the channel. Based upon the biological hydrophobicity scale, a significant number of these segments are predicted not to insert efficiently into a membrane due to their low hydrophobicity [126]. It was therefore a long-standing question how such H-segments insert [127,128]. One answer is that charged residues, especially arginine, have very favorable interactions with the phosphates of phospholipids [87,129].

Charged and polar residues can face a high energetic barrier when inserting into a lipid bilayer. Nevertheless, positive and negative charges within the same [130] and different TMHs [131] can interact




with each other, thereby drastically reducing this barrier. This concept can be extended to polar side chains, which are partially masked from the lipid bilayer by interacting with each other [56,132]. In fact, it was proposed early on that MPs are “inside-out” proteins, in the sense of having a polar core and a hydrophobic exterior, opposite to water-soluble proteins [133,134]. Although this idea is oversimplified, it suggests how the energy barrier for membrane insertion of hydrophilic side chains might be overcome and how specific tertiary structure interactions formed at the same time. It has been suggested that polar residues within TMHs can be used to predict the tertiary structure of polytopic MPs [126,135]. Individual TMHs can associate well after insertion into a membrane to form a functionally folded protein [11,12]. This implies that the specific interactions between TMHs that determine the final fold do not strictly require the action of a complex folding apparatus, as expected for equilibrium folding.

When, then, are specific interactions between TMHs formed? The simplistic and sequential insertion of TMHs one by one as discussed earlier implies that these interactions form after the TMHs have been inserted. While this might be the case for sufficiently hydrophobic TMHs, marginally hydrophobic segments would not insert into a membrane

efficiently; thus, more complex models of membrane insertion and folding are required. One could imagine that these interactions between TMHs occur early, before insertion into a membrane, either once a TMH exits the ribosome or once it enters the translocon. It has been shown that helical hairpins composed of two closely spaced TMHs can form in the ribosome exit vestibule as the TMHs of a voltage sensor domain exit the ribosome [136] (1 in Fig. 14). Although many TMHs risk aggregation when associating in an aqueous environment, the exit vestibule of the ribosome could form a confined compartment that promotes tertiary structure formation [137]. It is presently unclear whether the helix–helix interactions observed by Tu *et al.* [136] in the ribosome exit vestibule are native contacts or whether they represent a folding intermediate, although initial results suggest native contacts. These and other results provide a new perspective on co-translational folding of TMHs. It will be interesting to learn if other proteins use this early folding space.

Remarkably, the translocon can accommodate more than two nascent chain segments; thus, specific interactions could also form before the segments insert into a membrane [138]. In this case, helical hairpins could form as a minimal prefolding unit and bury hydrophilic interactions within the helix–helix interface before entering the lipid bilayer (2a and 2b in



**Fig. 14.** Folding spaces during MP structure formation. During nascent chain synthesis by the ribosome (brown), helical segments (shown in blue and orange) can associate already in the ribosomal vestibule (1) [136] or within the translocon channel (2a, green) [138], where polar residues can be shielded within the helix–helix interface (2b, red spheres) and hydrophobic side chains (orange spheres) form a membrane insertion competent surface. Helices can furthermore interact during insertion into a membrane (3, grey) whereas one segment is already inserted into the membrane and drives the insertion of a less hydrophobic segment by shielding hydrophilic residues within the interaction interface [143]. Interactions of helices within different proteins can also occur co-translationally (5) and was shown to drive MP complex assembly [149].

Fig. 14). It is unclear how the formation of such interactions is induced within the translocon, how water is expelled from the helix–helix interface, or if such early interactions are merely the result of tertiary interactions already formed in the ribosome's exit vestibule.

Currently, a widely discussed mode of the co-translational formation of helix–helix interactions is the formation during the sequential exit of successive TMHs through the lateral gate, where previously inserted TMHs that are in close vicinity of the lateral gate can conceivably mediate membrane insertion and tertiary structure folding at the same time (3 in Fig. 14). Sadlish *et al.* have shown that multiple TMHs of the same polypeptide chain can be associated with the eukaryotic translocon and mediate the efficient insertion of more C-terminally located TMHs [114]. This has also been confirmed for other MPs [139], suggesting that the translocon itself might be a foldase. However, it is unclear how and where TMHs associate with the exterior of the translocon complex. Although a site has been identified by crosslinking studies, it is unclear if the majority of TMHs occupy the same site [114]. Any specific interactions with a translocon complex that manages thousands of different TM segments would have to be replaced by stronger interactions with the protein's own TMHs as the newly inserted segments emerge from the complex. In such a scheme, a strict hierarchy of interaction strengths in the form of an interaction and folding blueprint would have to be in place for all MPs using the pathway. This might be achieved by the strict coupling of correct tertiary structure formation and membrane insertion. However, this blueprint would have to be at least partially encoded in the nascent chain itself, possibly in the form of conserved polar residues within TMHs [140]. If this hierarchical mechanism of tertiary structure formation is a common one, it will be interesting in the future to decipher the individual strengths of these interactions and thereby come closer to an *ab initio* understanding of MP folding. As pointed out above, other factors such as the point of secondary structure formation, tertiary structure formation, and retention within the translocon are critical factors as well.

An interesting special case is the folding of membrane channels that are composed of evolutionarily related halves [141,142]. Here, each half-channel is (or once was) a separate folding entity; otherwise, the high level of similarity between the halves could create interchangeability between helix–helix interactions from one half with the other that subsequently would result in protein misfolding.

In bacteria, the formation of specific interactions between side chains from interacting TMHs has been observed by means of APs (see above) [143] for several polytopic MPs during their insertion by the bacterial translocon. Mutating interacting residues in

either a membrane inserting segment or an already inserted segment in the same polypeptide chain decreased the pulling force detected by a C-terminally located AP. APs thus offer a highly sensitive *in vivo* tool for studying the co-translational formation of tertiary structure contacts, similar to *in vitro* AFM (see MP intrinsic interactions: TM helix folding and stability).

A recent publication offers unprecedented structural insight into the biogenesis of a polytopic MP in bacteria [144]. The structure of a ribosome-nascent chain-translocon complex shows the first two TMHs of proteorhodopsin (PR) inserted into the membrane just outside the lateral gate. Although the resolution of the structure is too low to reveal side-chain interactions between the PR TMHs themselves or with the translocon, an interaction of positively charged residues in the PR loop between TMH1 and TMH2 with the ribosomal RNA helix 59 could be observed. This interaction was verified by mutagenesis of the charged residues, which reduced the amount of co-purified SecY/nascent chain complex. Interestingly, the lateral gate was in between the fully open and closed conformation, which would allow a translocated hydrophilic segment to probe for tertiary interactions with previously inserted TM segments. It is tempting to speculate about possible mechanisms of membrane insertion and folding based on these results. Nevertheless, the structure represents, at best, a snapshot of the biogenesis of PR. Future investigations and more structural snapshots of the biogenesis of polytopic MPs might yield a more generalized and conclusive picture.

The insertion of more hydrophilic TMHs can occur at a later stage of assembly as well, which requires a larger structural reorganization of the maturing protein (4 in Fig. 14). This could be observed for several MPs [93,145–147]. In these cases, TMH3 is inserted in the opposite topology and TMH2 and TMH4 are initially not inserted into the membrane. Upon the reorientation of TMH3 at a later stage, TMH2 and TMH4 are inserted into the membrane and the water channel folds into its final functional structure. In order to form the tertiary structure of the protein, large-scale rearrangements involve induction of large tilts during folding, as well as topology reversals [148]. In a particularly impressive case, the topology of a bacterial MP was found to be in a flexible state until a specific C-terminal charged residue was synthesized [109].

These examples demonstrate that the formation of the tertiary structure that allows more hydrophilic TMHs to insert into a membrane can occur before, during, and after membrane insertion of the newly synthesized protein. It is currently unknown which is the preferred mode of folding in polytopic MPs because only a few examples have been investigated to date. In order to achieve a broader understanding and gain a more general view, it is necessary to follow

closely the folding of many more polytopic MPs in a high-throughput manner.

The concept of specific interactions between TMHs allowing more hydrophilic parts of a protein to insert into a membrane can also be extended to the quaternary structure, when it comes to the homo-oligomerization of the same subunit. Furthermore, interactions between different subunits of a newly synthesized MP complex can form during membrane insertion (hetero-oligomerization; 5 in Fig. 14) as was recently demonstrated for the subunits of the T cell receptor [149]. Feige and Hendershot found that a marginally hydrophobic TMH that is not assembled into a hetero-oligomeric T cell receptor is translocated into the ER lumen, rapidly recognized by the ER quality control machinery, and subsequently degraded. However, polar residues in already inserted, interacting subunits guided the membrane insertion and assembly of individual T cell receptor subunits and thereby prevented degradation. This marks an important step toward a mechanistic understanding of how MP complexes are assembled in cells. It will be both very challenging and interesting to understand mechanistically the assembly of complexes located in the ER membrane, complexes located in the mitochondrial inner membrane, and complexes located in the thylakoid membrane.

MP secondary and tertiary structures can form at different stages: in the ribosome exit tunnel, the ribosome exit vestibule [136], within the translocon channel [115–118], during membrane insertion [39,88], and within a membrane. Each of these environments is characterized by unique properties and thereby jointly offer an extended folding space for the many different MPs that are chaperoned by the co-translational MP synthesis and insertion complexes. After exploring basic principles of MP synthesis, membrane insertion, and folding using model MPs, it is now important to venture into the widely unexplored folding space of cellular MPs and to follow and understand mechanistically their synthesis within a cell. Not only are these misfolding pathways likely to offer new insights into the biogenesis of MPs, but they will provide us also with a more quantitative view of which modes of folding are the major ones and which represent the exceptions.

## Perspectives and Outlook

Where do we stand in our understanding of MP biogenesis in mid-2014? First, it is clear that our knowledge is much more advanced for the SecYEG/Sec61 translocons than for systems such as the mitochondrial TOM–TIM translocons, the peroxisomal PEX translocons, and the chloroplast TIC–TOC translocons (e.g., see Refs. [150–152]). The SecYEG-associated YidC translocon is also slowly

yielding its secrets. In recent years, the field has moved forward in a major way thanks to structural studies of the Sec and YidC translocons, and there are now many structures of Sec translocases in complex with ribosomes. In parallel, a biophysical description of the basic insertion and folding processes is emerging through the confrontation of thermodynamic and kinetic studies of highly simplified systems (peptide–bilayer interactions, *in vitro* folding of purified proteins, MD calculations) with quantitative *in vivo* studies of the kinds described above (“biological” hydrophobicity scales, co-translational force measurements). It is satisfying to see that we are finally moving beyond the simple cartoons (Fig. 1) to more detailed structure–function descriptions.

Still, many open questions remain, even for the Sec-type translocons. How are holo-translocons—both prokaryotic and eukaryotic—put together and how do the different components cooperate? It is currently unclear how the numerous accessory components of the holo-translocons are recruited and what the dynamics of complex assembly/disassembly are. It is also unknown how the bacterial translocons, YidC and SecYEG, cooperate to insert MPs. What are the precise roles of the lateral gate and the plug domain in SecYEG, what controls their movements, and how do they interact with polar and non-polar segments in a nascent chain? How strong is the functional coupling between the ribosome and the translocon; that is, can conformational changes resulting from interactions between a nascent chain and the ribosome be transmitted to the translocon?

Lipids appear to play an important role in protein translocation, but possible mechanisms are largely speculative (reviewed in Ref. [153]). From the point of view of a substrate protein, it is not clear to what extent a nascent chain can interact with membrane lipids as it passes from the ribosome into the translocon or at which point during translation that a TMH first starts to integrate into the bilayer. It seems clear that an incoming TMH can make specific interactions with already inserted TMHs at very early stages of membrane insertion, but how are the upstream TMHs “chaperoned” by the holo-translocon while waiting for downstream TMHs to appear? Further, what is the physical basis for the late reorientation of TMHs that has been seen with, for example, AQP1 [154], EmrE [109], and Band III [147], and when the membrane lipid composition is drastically changed [155]?

We tried to capture in Fig. 1 the idea deeply embedded in the literature that the nascent chain, powered by the GTP-based elongation energy of the ribosome, enters the translocon whereupon the hydrophobic ring opens, the plug domain moves out of the way, and the nascent chain passes through. If a suitable hydrophobic segment arrives, it is diverted into




the membrane by simple partitioning between translocon and bilayer [83,84,87,156]. Reflecting on the literature we have reviewed here, we wonder if this scheme is entirely correct. The translocon is absolutely essential for the assembly of most multi-span MPs, but there is a notable exception: the transmembrane protein KdpD.

KdpD, which acts as a potassium sensor in *E. coli* [59,157], has several features that are intriguing: The protein is tethered to the cytoplasmic surface of the inner membrane by four TM segments (residues 401–498) with very short interhelical loops that occur far downstream from the amino-terminus. It is targeted to the inner membrane by SRP recognition of a cytoplasmic amphipathic helix (residues 22–48) [158], and membrane insertion does not require SecA, SecE, or YidC [59]. Because KdpD does not apparently engage SecA or the SecYEG channel, a reasonable explanation is spontaneous insertion following targeting by the SRP to, presumably, the SRP receptor FtsY. In *E. coli*, FtsY is not permanently anchored to the inner membrane, but rather, it partitions between membrane and cytoplasm via an amphipathic helix [159]. This suggests that SRP recognizes the TMHs of KdpD and

targets them to FtsY on the membrane, which allows the TMHs to partition spontaneously. Such a scheme, while highly unusual, is completely consistent with what we know about lipid–protein interactions described at the outset in [Biophysical Boundaries of MP Insertion and Folding](#).

In order to be provocative, which was our charge from the editors of this issue, we propose an alternative view of the translocon-aided insertion of multi-span MPs and the secretion of soluble proteins (Fig. 15). Keeping in mind the huge thermodynamic driving forces for MP folding and assembly, we suggest that a TMH initially contacts the cytoplasmic membrane interface in the vicinity of the translocon and that it never fully enters the translocon channel but rather slides into the membrane along the lateral gate. This is not to say that the translocon cannot form a proper channel through the membrane, but we suggest that it does this only for secreted proteins and soluble domains or loop regions in MPs, as shown in Fig. 15b. The view is based upon four important observations. First, most hydrophobic and amphipathic helices have a very strong affinity for the interface region of lipid bilayers; it would be



**Fig. 15.** An alternative view of translocon-aided insertion of multi-span MPs and the secretion of soluble proteins. The idea of this alternative view is shown in a cartoon fashion in (a) (also see Fig. 10). We suggest that initial contact of the nascent chain (green, with red TMHs) is with the membrane interface in the vicinity of the translocon (blue) and that the chain does not immediately thread into the translocon. Rather, the translocon has YidC-like behavior; it provides a pathway for polar components of MPs to cross the membrane. This is not to say that it cannot form a passageway through the membrane, but we suggest that it does this only for polar polypeptide segments in MPs and for secreted proteins, as shown in (b).

thermodynamically surprising if nascent peptide chains of MPs did not interact with the membrane interface at some point during insertion. Second, so far, no structural data of active ribosome–translocon complexes reveal TM segments within the translocon channel; they are only seen at lateral gate exit site. Third, ribosome–translocon complexes are not apparently shielded from the cytoplasm and inner membrane surface; consistent with earlier observations, the recent high-resolution structure of a mammalian ribosome–Sec61 complex shows significant gap between the ribosome and the surface of the membrane [64]. Fourth, mitochondrial inner membranes in general do not have a SecY apparatus, only the YidC-equivalent Oxa1, suggesting that direct contact with the membrane is an essential feature of MP insertion and that nascent mitochondrial proteins insert and fold spontaneously into the membrane aided by Oxa1 to move charged and/or highly polar residues across the membrane.

An important consideration is that we do not know when or how nascent MP chains make initial contact with translocons or in what manner. Do they move directly from the exit tunnel straight into the translocon without intervening steps, as current cartoons suggest (Fig. 1)? Given relative target sizes, it seems possible that nascent chains could make first contact with the membrane in the vicinity of the translocon. This provides a logical explanation for the membrane insertion of the potassium channel S3b–S4 helices that apparently assemble in the ribosome vestibule [136]. Furthermore, it was shown recently that the ribosome detached from the translocon when a hydrophobic segment was placed in the ribosome exit tunnel and only hydrophilic stretches within the tunnel provided a strong attachment of the ribosome to the translocon [144]. The scheme would also explain why changes in *E. coli* inner membrane lipid composition affect the topology of LacY [155]. Consistent with the importance of thermodynamic driving forces in MP folding, Vitrac *et al.* have shown that so-called topology switching of LacY can occur *in vitro* in lipid vesicles [15].

We are attracted to the scheme depicted in Figs. 10b and 15a because of its simplicity: A TM segment initially associates with the top of the lateral gate and the surrounding membrane interface and strong thermodynamic forces then drive membrane partitioning. Such a process would be consistent with both crosslinking results and the AP force measurements (Fig. 13). It is not necessary in this scheme for the elongating chain to be threaded back and forth across the membrane as in Fig. 1. For the four-TMH protein shown in the Fig. 15a schematic, the translocon is mainly required for only two steps: the passage of the TM1–TM2 and TM3–TM4 connecting loops across the membrane, reminiscent of the helical hairpin insertion hypothesis [23]. Thermodynamic forces are sufficient for membrane partitioning, with the translocon acting as a catalyst.

Why do we need translocons at all if insertion and folding are driven by thermodynamics? The answer, of course, is that targeting and coordinated assembly of MPs are essential in the extraordinarily complicated environment of cells where thousands of biologically important reactions occur simultaneously in a crowded environment. Without translocons and all the other members of the MP assembly apparatus, chaos would prevail.

## Acknowledgements

This work was supported in part by grants from the National Institutes of General Medical Sciences (RO1 GM074637) and from the National Institute of General Medical Sciences and the National Institute of Neurological Disorders and Stroke jointly (PO1 GM86685) to S.H.W.; by grants from the Swedish Cancer Foundation, the Swedish Research Council, the Swedish Foundation for Strategic Research, the European Research Council (ERC-2008-AdG 232648), and the Knut and Alice Wallenberg Foundation to G.v.H.; and by a research fellowship from the Deutsche Forschungsgemeinschaft (CY 74/1-1) to F.C.

## Q&A with Stephen White and Gunnar von Heijne

### Q1

**FJD:** *In your rather provocative model a TM span does not enter the translocon but slides along the lateral gate. How can this model explain that some mutations of residues in the constriction ring of the translocon have an impact on TM span integration?*

**FC/GvH/SW:** The crucial issues are (1) the time scale of the integration and (2) the dynamics of the translocon. Presumably, integration of TM segments occurs on the same time scale as the elongation rate of nascent polypeptide chains, which is roughly one amino acid per 50 msec or about 1 second for a single TM segment or signal sequence. We do not know what conformational states are available to a translocon interacting with a nascent chain, but we suggest that on such long time scales the nascent chain has ample opportunity to explore rather thoroughly the surrounding lipid bilayer and interior of the translocon, including the so-called constriction ring. The standard view that the nascent chain passes through the center of the translocon and exits through the lateral gate implies that the reverse process is also possible. The slow

time scale likely permits the chain to explore all of the states energetically available to it, including ones separated by significant kinetic barriers. Thus, we suggest that even if the primary entrance path of the chain is via the lateral gate, the amino acid composition of the constriction ring matters in the fate of the nascent chain. What is missing in our understanding of the role of the passive translocon apparatus is knowledge of the conformational states available to the translocon and the energy barriers that separate the states. Molecular dynamics simulations could help clarify conformational states available; unfortunately times scales of tens of milliseconds are not technically feasible at present.

## Q2

**FJD:** *How does your model account for the positive inside rule for integration of membrane proteins with a type II topology? In your model Sec61 should not be sensitive to the N-terminal flanking segment, while genetic evidence indicates that point mutations in Sec61 can affect the topology of reporter proteins.*

**FC/GvH/SW:** Our response to question 1 above applies in this case as well. Sec61/SecY mutations that affect topology presumably affect translocon/nascent-chain conformational states and energy landscape. In simplistic terms, the thermodynamically favored location of a translocating chain segment in the translocon may be affected by its hydrophobicity: strongly hydrophobic segments tend to translocate (or insert into the membrane, as the case may be) in a more lipid-exposed location on the outside of the lateral gate, while less hydrophobic segments may tend to be more buried within the translocon channel as they move through the membrane. Topology-determining charged residues that flank a hydrophobic transmembrane segment could therefore tend to be located more deeply in the lateral gate, or even fully within the translocon channel, when in transit. As they exit the ribosome, they may also transiently be exposed to lipid headgroups.

## Q3

**FJD:** *According to your model, the signal sequence or first TM span of a membrane protein positions the nascent chain in the vicinity of the translocon, and the C-terminal flanking segment slides into the lateral gate in a hairpin conformation. How does a protein segment from the mature region promote channel opening?*

**FC/GvH/SW:** Our view of the translocon as a dynamic molecule with multiple conformational states

(response 1) eliminates the need to think of TM helix insertion as series of discrete steps involving distinct opening and closing events. We suggest that integration is a continuous process in which the growing nascent chain persistently explores the states available to it. We suppose that translocon mutations (response 2) affect these states and the kinetic barriers that separate them.

Received 25 June 2014;

Received in revised form 13 September 2014;

Accepted 22 September 2014

Available online 30 September 2014

## Keywords:

membrane protein folding;  
membrane protein biogenesis;  
transmembrane helix;  
lipid-protein interactions

†F.C., G.v.H., and S.H.W. contributed equally to this work and are listed alphabetically.

## Abbreviations used:

TM, transmembrane; TMH, transmembrane helix; AP, arrest peptide; bR, bacteriorhodopsin; MP, membrane protein; ER, endoplasmic reticulum; MD, molecular dynamics; AFM, atomic force microscopy; PTC, polypeptide transferase center; SRP, signal recognition particle; PR, proteorhodopsin.

## References

- [1] White SH, Wimley WC. Membrane protein folding and stability: Physical principles. *Annu Rev Biophys Biomol Struct* 1999;28:319–65.
- [2] Popot JL, Engelman DM. Helical membrane protein folding, stability, and evolution. *Annu Rev Biochem* 2000;69:881–922.
- [3] MacKenzie KR. Folding and stability of  $\alpha$ -helical integral membrane proteins. *Chem Rev* 2006;106:1931–77.
- [4] Goldenberg DP. Mutational analysis of protein folding and stability. In: Creighton TE, editor. *Protein Folding*. New York: W. H. Freeman; 1992. p. 353–403.
- [5] Yang AS, Sharp KA, Honig B. Analysis of the heat capacity dependence of protein folding. *J Mol Biol* 1992;227:889–900.
- [6] Hong H, Joh NH, Bowie JU, Tamm LK. Methods for measuring the thermodynamic stability of membrane proteins. *Methods Enzymol* 2009;455:213–36.
- [7] Jackson MB, Sturtevant JM. Phase transitions of the purple membranes of *Halobacterium halobium*. *Biochemistry* 1978;17:911–5.
- [8] Lau FW, Bowie JU. A method for assessing the stability of a membrane protein. *Biochemistry* 1997;36:5884–92.
- [9] Popot J-L, Gerchman S-E, Engelman DM. Refolding of bacteriorhodopsin in lipid bilayers. A thermodynamically controlled two-stage process. *J Mol Biol* 1987;198:655–76.
- [10] Popot JL, Trewheila J, Engelman DM. Reformation of crystalline purple membrane from purified bacteriorhodopsin fragments. *EMBO J* 1986;5:3039–44.



- [11] Ridge KD, Lee SSJ, Yao LL. *In vivo* assembly of rhodopsin from expressed polypeptide fragments. *Proc Natl Acad Sci U S A* 1995;92:3204–8.
- [12] Bibi E, Kaback H. *In vivo* expression of the *lacY* gene in two segments leads to functional *lac* permease. *Proc Natl Acad Sci U S A* 1990;87:4325–9.
- [13] Groves JD, Tanner MJA. Topology studies with biosynthetic fragments identify interacting transmembrane regions of the human red-cell anion exchanger (band 3; AE1). *Biochem J* 1999;344:687–97.
- [14] Koebnik R. *In vivo* membrane assembly of split variants of the *E. coli* outer membrane protein OmpA. *EMBO J* 1996;15:3529–37.
- [15] Vitrac H, Bogdanov M, Dowhan W. *In vitro* reconstitution of lipid-dependent dual topology and post-assembly topological switching of a membrane protein. *Proc Natl Acad Sci U S A* 2013;110:9338–43.
- [16] Bayburt TH, Sligar SG. Membrane protein assembly into nanodiscs. *FEBS Lett* 2010;584:1721–7.
- [17] Kalmbach R, Chizhov I, Schumacher MC, Friedrich T, Bamberg E, Engelhard M. Functional cell-free synthesis of a seven helix membrane protein: *In situ* insertion of bacteriorhodopsin into liposomes. *J Mol Biol* 2007;371:639–48.
- [18] Goren MA, Nozawa A, Makino SI, Wrobel RL, Fox BG. Cell-free translation of integral membrane proteins in unilamellar liposomes. *Methods Enzymol* 2009;463:647–73.
- [19] Long AR, O'Brian CC, Alder NN. The cell-free integration of a polytopic mitochondrial membrane protein into liposomes occurs cotranslationally and in a lipid-dependent manner. *PLoS One* 2012;7:e46332.
- [20] Roos C, Zocher M, Müller D, Münch D, Schneider T, Sahl HG, et al. Characterization of co-translationally formed nanodisc complexes with small multidrug transporters, proteorhodopsin, and with the *E. coli* MraY translocase. *Biochim Biophys Acta* 2012;1818:3098–106.
- [21] Yildiz AA, Knoll W, Gennis RB, Sinner EK. Cell-free synthesis of cytochrome *bo3* ubiquinol oxidase in artificial membranes. *Anal Biochem* 2012;423:39–45.
- [22] Gloge F, Becker AH, Kramer G, Bukau B. Co-translational mechanisms of protein maturation. *Curr Opin Struct Biol* 2014;24:24–33.
- [23] Engelman DM, Steitz TA. On the folding and insertion of globular membrane proteins. In: Wetlaufer DB, editor. *The Protein Folding Problem*. Boulder: Westview Press; 1984. p. 87–113.
- [24] Wimley WC, Creamer TP, White SH. Solvation energies of amino acid sidechains and backbone in a family of host-guest pentapeptides. *Biochemistry* 1996;35:5109–24.
- [25] Wimley WC, White SH. Experimentally determined hydrophobicity scale for proteins at membrane interfaces. *Nat Struct Biol* 1996;3:842–8.
- [26] Wimley WC, Gawrisch K, Creamer TP, White SH. A direct measurement of salt-bridge solvation energies using a peptide model system: Implications for protein stability. *Proc Natl Acad Sci U S A* 1996;93:2985–90.
- [27] Reynolds JA, Gilbert DB, Tanford C. Empirical correlation between hydrophobic free energy and aqueous cavity surface area. *Proc Natl Acad Sci U S A* 1974;71:2925–7.
- [28] Wimley WC, Hristova K, Ladokhin AS, Silvestro L, Axelsen PH, White SH. Folding of beta-sheet membrane proteins: A hydrophobic hexapeptide model. *J Mol Biol* 1998;277:1091–110.
- [29] Ladokhin AS, White SH. Folding of amphipathic alpha-helices on membranes: Energetics of helix formation by melittin. *J Mol Biol* 1999;285:1363–9.
- [30] Ben-Tal N, Sitkoff D, Topol IA, Yang AS, Burt SK, Honig B. Free energy of amide hydrogen bond formation in vacuum, in water, and in liquid alkane solution. *J Phys Chem B* 1997;101:450–7.
- [31] Ben-Tal N, Ben-Shaul A, Nicholls A, Honig B. Free energy determinants of alpha-helix insertion into lipid bilayers. *Biophys J* 1996;70:1803–12.
- [32] Almeida PF, Ladokhin AS, White SH. Hydrogen-bond energetics drive helix formation in membrane interfaces. *Biochim Biophys Acta* 2012;1818:178–82.
- [33] Jayasinghe S, Hristova K, White SH. Energetics, stability, and prediction of transmembrane helices. *J Mol Biol* 2001;312:927–34.
- [34] Segrest JP, Jackson RL, Marchesi VT, Guyer RB, Terry W. Red cell membrane glycoprotein: Amino acid sequence of an intramembranous region. *Biochem Biophys Res Commun* 1972;49:964–9.
- [35] Sapra K, Besir S, Oesterhelt D, Muller D. Characterizing molecular interactions in different bacteriorhodopsin assemblies by single-molecule force spectroscopy. *J Mol Biol* 2006;355:640–50.
- [36] Oesterhelt F, Oesterhelt D, Pfeiffer M, Engel A, Gaub HE, Muller DJ. Unfolding pathways of individual bacteriorhodopsins. *Science* 2000;288:143–6.
- [37] Jarzynski C. Equilibrium free-energy differences from non-equilibrium measurements: A master-equation approach. *Phys Rev E* 1997;56:5018–35.
- [38] Müller DJ, Kessler M, Oesterhelt F, Möller C, Oesterhelt D, Gaub H. Stability of bacteriorhodopsin  $\alpha$ -helices and loops analyzed by single-molecule force spectroscopy. *Biophys J* 2002;83:3578–88.
- [39] Ulmschneider JP, Smith JC, White SH, Ulmschneider MB. *In silico* partitioning and transmembrane insertion of hydrophobic peptides under equilibrium conditions. *J Am Chem Soc* 2011;133:15487–95.
- [40] Andersson M, Ulmschneider JP, Ulmschneider MB, White SH. Conformational states of melittin at a bilayer interface. *Biophys J* 2013;104:L12–4.
- [41] Chen CH, Wiedman G, Khan A, Ulmschneider MB. Absorption and folding of melittin onto lipid bilayer membranes via unbiased atomic detail microsecond molecular dynamics simulation. *Biochem Biophys Acta* 2014;1838:2243–9.
- [42] Popot JL, Engelman DM. Membrane protein folding and oligomerization: The 2-stage model. *Biochemistry* 1990;29:4031–7.
- [43] Furthmayr H, Marchesi VT. Subunit structure of human erythrocyte glycophorin A. *Biochemistry* 1976;15:1137–44.
- [44] Bormann B-J, Knowles WJ, Marchesi VT. Synthetic peptides mimic the assembly of transmembrane glycoproteins. *J Biol Chem* 1989;264:4033–7.
- [45] Lemmon MA, Flanagan JM, Treutlein HR, Zhang J, Engelman DM. Sequence specificity in the dimerization of transmembrane  $\alpha$ -helices. *Biochemistry* 1992;31:12719–25.
- [46] Lemmon MA, Flanagan JM, Hunt JF, Adair BD, Bormann B-J, Dempsey CE, et al. Glycophorin A dimerization is driven by specific interactions between transmembrane  $\alpha$ -helices. *J Biol Chem* 1992;267:7683–9.
- [47] MacKenzie KR, Prestegard JH, Engelman DM. A transmembrane helix dimer: Structure and implications. *Science* 1997;276:131–3.
- [48] Langosch D, Brosig B, Kolmar H, Fritz HJ. Dimerisation of the glycophorin A transmembrane segment in membranes

- probed with the ToxR transcription activator. *J Mol Biol* 1996;263:525–30.
- [49] Eriksson AE, Baase WA, Zhang XJ, Heinz DW, Blaber M, Baldwin EP, et al. Response of a protein structure to cavity-creating mutations and its relation to the hydrophobic effect. *Science* 1992;255:178–83.
  - [50] Lemmon MA, Engelman DM. Specificity and promiscuity in membrane helix interactions. *Q Rev Biophys* 1994;27:157–218.
  - [51] Russ WP, Engelman DM. The GxxxG motif: A framework for transmembrane helix–helix association. *J Mol Biol* 2000;296:911–9.
  - [52] Senes A, Gerstein M, Engelman DM. Statistical analysis of amino acid patterns in transmembrane helices: The GxxxG motif occurs frequently and in association with beta-branched residues at neighboring positions. *J Mol Biol* 2000;296:921–36.
  - [53] Doura AK, Fleming KG. Complex interactions at the helix–helix interface stabilize the glycophorin A transmembrane dimer. *J Mol Biol* 2004;343:1487–97.
  - [54] Hong H, Bowie JU. Dramatic destabilization of transmembrane helix interactions by features of natural membrane environments. *J Am Chem Soc* 2011;133:11389–98.
  - [55] Chen L, Novicky L, Merzlyakov M, Hristov T, Hristova K. Measuring the energetics of membrane protein dimerization in mammalian membranes. *J Am Chem Soc* 2010;132:3628–35.
  - [56] Zhou FX, Merianos HJ, Brunker AT, Engelman DM. Polar residues drive association of polyleucine transmembrane helices. *Proc Natl Acad Sci U S A* 2001;98:2250–5.
  - [57] Hong H, Blois T, Cao Z, Bowie JU. Method to measure strong protein–protein interactions in lipid bilayers using a steric trap. *Proc Natl Acad Sci U S A* 2010;107:19802–36.
  - [58] Chang Y-C, Bowie JU. Measuring membrane protein stability under native conditions. *Proc Natl Acad Sci U S A* 2014;111:219–24.
  - [59] Facey SJ, Kuhn A. The sensor protein KdpD inserts into the *Escherichia coli* membrane independent of the Sec translocase and YidC. *Eur J Biochem* 2003;270:1724–34.
  - [60] Cech TR. Structural biology. The ribosome is a ribozyme. *Science* 2000;289:878–9.
  - [61] Ban N, Nissen P, Hansen J, Moore PB, Steitz TA. The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science* 2000;289:905–20.
  - [62] Ben-Shem A, Garreau de Loubresse N, Melnikov S, Jenner L, Yusupova G, Yusupov M. The structure of the eukaryotic ribosome at 3.0 Å resolution. *Science* 2011;334:1524–9.
  - [63] Armache JP, Jarasch A, Anger AM, Villa E, Becker T, Bhushan S, et al. Cryo-EM structure and rRNA model of a translating eukaryotic 80S ribosome at 5.5-Å resolution. *Proc Natl Acad Sci U S A* 2010;107:19748–53.
  - [64] Voorhees RM, Fernández IS, Scheres SHW, Hegde RS. Structure of the mammalian ribosome–Sec61 complex to 3.4 Å resolution. *Cell* 2014;157:1632–43.
  - [65] Keenan RJ, Freymann DM, Stroud RM, Walter P. The signal recognition particle. *Annu Rev Biochem* 2001;70:755–75.
  - [66] Akopian D, Shen K, Zhang X, Shan SO. Signal recognition particle: An essential protein-targeting machine. *Annu Rev Biochem* 2013;82:693–721.
  - [67] Flanagan JJ, Chen JC, Miao Y, Shao Y, Lin J, Bock PE, et al. Signal recognition particle binds to ribosome-bound signal sequences with fluorescence-detected subnanomolar affinity that does not diminish as the nascent chain lengthens. *J Biol Chem* 2003;278:18628–37.
  - [68] Ataíde SF, Schmitz N, Shen K, Ke A, Shan SO, Doudna JA, et al. The crystal structure of the signal recognition particle in complex with its receptor. *Science* 2011;331:881–6.
  - [69] Saraogi I, Akopian D, Shan SO. A tale of two GTPases in cotranslational protein targeting. *Protein Sci* 2011;20:1790–5.
  - [70] Janda CY, Li J, Oubridge C, Hernandez H, Robinson CV, Nagai K. Recognition of a signal peptide by the signal recognition particle. *Nature* 2010;465:507–10.
  - [71] Ito K, Chiba S, Pogliano K. Divergent stalling sequences sense and control cellular physiology. *Biochem Biophys Res Commun* 2010;393:1–5.
  - [72] Ramu H, Mankin A, Vazquez-Laslop N. Programmed drug-dependent ribosome stalling. *Mol Microbiol* 2009;71:811–24.
  - [73] Drew D, Fröderberg L, Baars L, de Gier J-WL. Assembly and overexpression of membrane proteins in *Escherichia coli*. *Biochim Biophys Acta* 2003;1610:3–10.
  - [74] Dalbey RE, Kuhn A, Zhu L, Kiefer D. The membrane insertase YidC. *Biochim Biophys Acta* 1843;2014:1489–96.
  - [75] Denks K, Vogt A, Sachelaru I, Petriman NA, Kudva R, Koch HG. The Sec translocon mediated protein transport in prokaryotes and eukaryotes. *Mol Membr Biol* 2014;31:58–84.
  - [76] Greber BJ, Boehringer D, Leitner A, Bieri P, Voigts-Hoffmann F, Erzberger JP, et al. Architecture of the large subunit of the mammalian mitochondrial ribosome. *Nature* 2014;505:515.
  - [77] Rapoport TA. Protein translocation across the eukaryotic endoplasmic reticulum and bacterial plasma membranes. *Nature* 2007;450:663–9.
  - [78] van den Berg B, Clemons WM, Collinson I, Modis Y, Hartmann E, Harrison SC, et al. X-ray structure of a protein-conducting channel. *Nature* 2004;427:36–44.
  - [79] Tsukazaki T, Mori H, Fukai S, Ishitani R, Mori T, Dohmae N, et al. Conformational transition of Sec machinery inferred from bacterial SecYE structures. *Nature* 2008;455:988–91.
  - [80] Egea PF, Stroud RM. Lateral opening of a translocon upon entry of protein suggests the mechanism of insertion into membranes. *Proc Natl Acad Sci U S A* 2010;107:17182–7.
  - [81] Kihara A, Akiyama Y, Ito K. FtsH is required for proteolytic elimination of uncomplexed forms of SecY, an essential protein translocase subunit. *Proc Natl Acad Sci U S A* 1995;92:4532–6.
  - [82] van der Sluis EO, van der Vries E, Berrelkamp G, Nouwen N, Driessen AJ. Topologically fixed SecE is fully functional. *J Bacteriol* 2006;188:1188–90.
  - [83] Hessa T, Meindl-Beinker NM, Bernsel A, Kim H, Sato Y, Lerch-Bader M, et al. Molecular code for transmembrane-helix recognition by the Sec61 translocon. *Nature* 2007;450:1026–30.
  - [84] Hessa T, Kim H, Bihlmaier K, Lundin C, Boekel J, Andersson H, et al. Recognition of transmembrane helices by the endoplasmic reticulum translocon. *Nature* 2005;433:377–81.
  - [85] Öjemalm K, Botelho SC, Studle C, von Heijne G. Quantitative analysis of SecYEG-mediated insertion of transmembrane alpha-helices into the bacterial inner membrane. *J Mol Biol* 2013;425:2813–22.
  - [86] Öjemalm K, Higuchi T, Jiang Y, Langel U, Nilsson I, White SH, et al. Apolar surface area determines the efficiency of translocon-mediated membrane-protein integration into the endoplasmic reticulum. *Proc Natl Acad Sci U S A* 2011;108:E359–64.
  - [87] Schow EV, Freites JA, Cheng P, Bernsel A, von Heijne G, White SH, et al. Arginine in membranes: The connection

- between molecular dynamics simulations and translocon-mediated insertion experiments. *J Membr Biol* 2011;239:35–48.
- [88] Ulmschneider MB, Ulmschneider JP, Schiller N, Wallace BA, von Heijne G, White SH. Spontaneous transmembrane helix insertion thermodynamically mimics translocon-guided insertion. *Nat Commun* 2014;5:4863.
- [89] Gogala M, Becker T, Beatrix B, Armache J-P, Barrio-Garcia C, Berninghausen O, et al. Cryo-EM structures of the Sec61-complex engaged in nascent peptide translocation or membrane insertion. *Nature* 2014;506:107–10.
- [90] Park E, Menetret JF, Gumbart JC, Ludtke SJ, Li W, Whynot A, et al. Structure of the SecY channel during initiation of protein translocation. *Nature* 2013;506:102–6.
- [91] Zhang B, Miller TF. Long-timescale dynamics and regulation of Sec-facilitated protein translocation. *Cell Rep* 2012;2:927–37.
- [92] Mackinnon AL, Paavilainen VO, Sharma A, Hegde RS, Taunton J. An allosteric Sec61 inhibitor traps nascent transmembrane helices at the lateral gate. *eLife* 2014;3:e01483.
- [93] Kauko A, Hedin LE, Thebaud E, Cristobal S, Elofsson A, von Heijne G. Repositioning of transmembrane  $\alpha$ -helices during membrane protein folding. *J Mol Biol* 2010;397:190–201.
- [94] Hedin LE, Öjemalm K, Bernsel A, Hennerdal A, Illergård K, Enquist K, et al. Membrane insertion of marginally hydrophobic transmembrane helices depends on sequence context. *J Mol Biol* 2010;396:221–9.
- [95] Öjemalm K, Halling KK, Nilsson I, von Heijne G. Orientational preferences of neighboring helices can drive ER insertion of a marginally hydrophobic transmembrane helix. *Mol Cell* 2012;45:529–40.
- [96] Kumazaki K, Chiba S, Takemoto M, Furukawa A, Nishiyama K, Sugano Y, et al. Structural basis of Sec-independent membrane protein insertion by YidC. *Nature* 2014;509:516–20.
- [97] Wickles S, Singharoy A, Andreani J, Seemayer S, Bischoff L, Berninghausen O, et al. A structural model of the active ribosome-bound membrane protein insertase YidC. *eLife* 2014. <http://dx.doi.org/10.7554/eLife.03035>.
- [98] Lotz M, Haase W, Kuhlbrandt W, Collinson I. Projection structure of YidC: A conserved mediator of membrane protein assembly. *J Mol Biol* 2008;375:901–7.
- [99] Kohler R, Boehringer D, Greber B, Bingel-Erlenmeyer R, Collinson I, Schaffitzel C, et al. YidC and Oxa1 form dimeric insertion pores on the translating ribosome. *Mol Cell* 2009;34:344–53.
- [100] Kedrov A, Sustarsic M, de Keyser J, Caumanns JJ, Wu ZC, Driessen AJ. Elucidating the native architecture of the YidC:ribosome complex. *J Mol Biol* 2013;425:4112–24.
- [101] Seitzl I, Wickles S, Beckmann R, Kuhn A, Kiefer D. The C-terminal regions of YidC from *Rhodospirella baltica* and *Oceanicaulis alexandrii* bind to ribosomes and partially substitute for SRP receptor function in *Escherichia coli*. *Mol Microbiol* 2014;91:408–21.
- [102] Xie K, Hessa T, Seppälä S, Rapp M, von Heijne G, Dalbey RE. Features of transmembrane segments that promote the lateral release from the translocase into the lipid phase. *Biochemistry* 2007;46:15153–61.
- [103] Martoglio B, Hofmann MW, Brunner J, Dobberstein B. The protein-conducting channel in the membrane of the endoplasmic reticulum is open laterally toward the lipid bilayer. *Cell* 1995;81:207–14.
- [104] Butkus ME, Prundeanu LB, Oliver DB. Translocon “pulling” of nascent SecM controls the duration of its translational pause and secretion-responsive secA regulation. *J Bacteriol* 2003;185:6719–22.
- [105] Yap MN, Bernstein HD. The plasticity of a translation arrest motif yields insights into nascent polypeptide recognition inside the ribosome tunnel. *Mol Cell* 2009;34:201–11.
- [106] Ismail N, Hedman R, Schiller N, von Heijne G. A biphasic pulling force acts on transmembrane helices during translocon-mediated membrane integration. *Nat Struct Mol Biol* 2012;19:1018–22.
- [107] von Heijne G. The distribution of positively charged residues in bacterial inner membrane proteins correlates with the trans-membrane topology. *EMBO J* 1986;5:3021–7.
- [108] Gafvelin G, von Heijne G. Topological “frustration” in multi-spanning *E. coli* inner membrane proteins. *Cell* 1994;77:401–12.
- [109] Seppälä S, Slusky JS, Lloris-Garcera P, Rapp M, von Heijne G. Control of membrane protein topology by a single C-terminal residue. *Science* 2010;328:1698–700.
- [110] Fujita H, Kida Y, Hagiwara M, Morimoto F, Sakaguchi M. Positive charges of translocating polypeptide chain retrieve an upstream marginal hydrophobic segment from the endoplasmic reticulum lumen to the translocon. *Mol Biol Cell* 2010;21:2045–56.
- [111] Demirci E, Junne T, Baday S, Berneche S, Spiess M. Functional asymmetry within the Sec61p translocon. *Proc Natl Acad Sci U S A* 2013;110:18856–61.
- [112] Junne T, Kocik L, Spiess M. The hydrophobic core of the Sec61 translocon defines the hydrophobicity threshold for membrane integration. *Mol Biol Cell* 2010;21:1662–70.
- [113] Pitonzo D, Yang Z, Matsumura Y, Johnson AE, Skach WR. Sequence-specific retention and regulated integration of a nascent membrane protein by the endoplasmic reticulum Sec61 translocon. *Mol Biol Cell* 2009;20:685–98.
- [114] Sadlish H, Pitonzo D, Johnson AE, Skach WR. Sequential triage of transmembrane segments by Sec61a during biogenesis of a native multispanning membrane protein. *Nat Struct Mol Biol* 2005;12:870–8.
- [115] Lu J, Deutsch C. Secondary structure formation of a transmembrane segment in Kv channels. *Biochemistry* 2005;44:8230–43.
- [116] Lu J, Deutsch C. Folding zones inside the ribosomal exit tunnel. *Nat Struct Mol Biol* 2005;12:1123–9.
- [117] Woolhead CA, Johnson AE, Bernstein HD. Translation arrest requires two-way communication between a nascent polypeptide and the ribosome. *Mol Cell* 2006;22:587–98.
- [118] Daniel CJ, Conti B, Johnson AE, Skach WR. Control of translocation through the Sec61 translocon by nascent polypeptide structure within the ribosome. *J Biol Chem* 2008;283:20864–73.
- [119] Kim S, Jeon TJ, Oberai A, Yang D, Schmidt JJ, Bowie JU. Transmembrane glycine zippers: Physiological and pathological roles in membrane proteins. *Proc Natl Acad Sci U S A* 2005;102:14278–83.
- [120] Fleming KG, Ackerman AL, Engelman DM. The effect of point mutations on the free energy of transmembrane  $\alpha$ -helix dimerization. *J Mol Biol* 1997;272:266–75.
- [121] Melnyk RA, Kim S, Curran AR, Engelman DM, Bowie JU, Deber CM. The affinity of GXXXG motifs in transmembrane helix-helix interactions is modulated by long-range communication. *J Biol Chem* 2004;279:16591–7.
- [122] Schneider D, Engelman DM. Motifs of two small residues can assist but are not sufficient to mediate transmembrane helix interactions. *J Mol Biol* 2004;343:799–804.

- [123] Liu Y, Engelman DM, Gerstein M. Genomic analysis of membrane protein families: Abundance and conserved motifs. *Genome Biol* 2002;3:0054 research.
- [124] Li E, Wimley WC, Hristova K. Transmembrane helix dimerization: Beyond the search for sequence motifs. *Biochim Biophys Acta* 1818;2012:183–93.
- [125] Fleishman SJ, Schlessinger J, Ben-Tal N. A putative molecular-activation switch in the transmembrane domain of erbB2. *Proc Natl Acad Sci U S A* 2002;99:15937–40.
- [126] Elofsson A, von Heijne G. Membrane protein structure: Prediction *versus* reality. *Annu Rev Biochem* 2007;76:125–40.
- [127] Bowie JU. Solving the membrane protein folding problem. *Nature* 2005;438:581–9.
- [128] von Heijne G. Formation of transmembrane helices *in vivo*—Is hydrophobicity all that matters? *J Gen Physiol* 2007;129:353–6.
- [129] Freites JA, Tobias DJ, von Heijne G, White SH. Interface connections of a transmembrane voltage sensor. *Proc Natl Acad Sci U S A* 2005;102:15059–64.
- [130] Chin CN, von Heijne G. Charge pair interactions in a model transmembrane helix in the ER membrane. *J Mol Biol* 2000;303:1–5.
- [131] Johnson ET, Parson WW. Electrostatic interactions in an integral membrane protein. *Biochemistry* 2002;41:6483–94.
- [132] Gratkowski H, Lear JD, DeGrado WF. Polar side chains drive the association of model transmembrane peptides. *Proc Natl Acad Sci U S A* 2001;98:880–5.
- [133] Rees DC, DeAntonio L, Eisenberg D. Hydrophobic organization of membrane proteins. *Science* 1989;245:510–3.
- [134] Stevens TJ, Arkin IT. Are membrane proteins “inside-out” proteins? *Proteins Struct Funct Genet* 1999;36:135–43.
- [135] Koehler J, Woetzel N, Staritzbichler R, Sanders CR, Meiler J. A unified hydrophobicity scale for multispan membrane proteins. *Proteins* 2009;76:13–29.
- [136] Tu L, Khanna P, Deutsch C. Transmembrane segments form tertiary hairpins in the folding vestibule of the ribosome. *J Mol Biol* 2014;426:185–98.
- [137] O'Brien EP, Christodoulou J, Vendruscolo M, Dobson CM. New scenarios of protein folding can occur on the ribosome. *J Am Chem Soc* 2011;133:513–26.
- [138] Zhang L, Sato Y, Hessa T, von Heijne G, Lee JK, Kodama I, et al. Contribution of hydrophobic and electrostatic interactions to the membrane integration of the Shaker K<sup>+</sup> channel voltage sensor domain. *Proc Natl Acad Sci U S A* 2007;104:8263–8.
- [139] Cross BC, High S. Dissecting the physiological role of selective transmembrane-segment retention at the ER translocon. *J Cell Sci* 2009;122:1768–77.
- [140] Illergård K, Kauko A, Elofsson A. Why are polar residues within the membrane core evolutionarily conserved? *Proteins* 2011;79:79–91.
- [141] Lloris-Garcera P, Seppala S, Slusky JS, Rapp M, von Heijne G. Why have small multidrug resistance proteins not evolved into fused, internally duplicated structures? *J Mol Biol* 2014;426:2246–54.
- [142] Duran AM, Meiler J. Inverted topologies in membrane proteins: A mini-review. *Comput Struct Biotechnol J* 2013;8:e201308004.
- [143] Cymer F, von Heijne G. Cotranslational folding of membrane proteins probed by arrest-peptide-mediated force measurements. *Proc Natl Acad Sci U S A* 2013;110:14640–5.
- [144] Bischoff L, Wickles S, Berninghausen O, van der Sluis EO, Beckmann R. Visualization of a polytopic membrane protein during SecY-mediated membrane insertion. *Nat Commun* 2014;5:4103.
- [145] Lu Y, Turnbull IR, Bragin A, Carveth K, Verkman AS, Skach WR. Reorientation of aquaporin-1 topology during maturation in the endoplasmic reticulum. *Mol Biol Cell* 2000;11:2973–85.
- [146] Watson HR, Wunderley L, Andreou T, Warwicker J, High S. Reorientation of the first signal-anchor sequence during potassium channel biogenesis at the Sec61 complex. *Biochem J* 2013;456:297–309.
- [147] Kanki T, Sakaguchi M, Kitamura A, Sato T, Mihara K, Hamasaki N. The tenth membrane region of band 3 is initially exposed to the luminal side of the endoplasmic reticulum and then integrated into a partially folded band 3 intermediate. *Biochemistry* 2002;41:13973–81.
- [148] Virkki M, Boekel C, Illergård K, Peters C, Shu N, Tsigiris KD, et al. Large tilts in transmembrane helices can be induced during tertiary structure formation. *J Mol Biol* 2014;426:2529–38.
- [149] Feige MJ, Hendershot LM. Quality control of integral membrane proteins by assembly-dependent membrane integration. *Mol Cell* 2013;51:297–309.
- [150] Rucktaschel R, Girzalsky W, Erdmann R. Protein import machineries of peroxisomes. *Biochim Biophys Acta* 1808;2011:892–900.
- [151] Sokol AM, Sztolsztener ME, Wasilewski M, Heinz E, Chacinska A. Mitochondrial protein translocases for survival and wellbeing. *FEBS Lett* 2014;588:2484–95.
- [152] Benz JP, Soll J, Bolter B. Protein transport in organelles: The composition, function and regulation of the Tic complex in chloroplast protein import. *FEBS J* 2009;276:1166–76.
- [153] van Voorst F, de Kruijff B. Role of lipids in the translocation of proteins across membranes. *Biochem J* 2000;347:601–12.
- [154] Buck TM, Wagner J, Grund S, Skach WR. A novel tripartite motif involved in aquaporin topogenesis, monomer folding and tetramerization. *Nat Struct Mol Biol* 2007;14:762–9.
- [155] Dowhan W, Bogdanov M. Lipid-dependent membrane protein topogenesis. *Annu Rev Biochem* 2009;78:515–40.
- [156] Heinrich S, Mothes W, Brunner J, Rapoport T. The Sec61p complex mediates the integration of a membrane protein by allowing lipid partitioning of the transmembrane domain. *Cell* 2000;102:233–44.
- [157] Zimmann P, Puppe W, Altendorf K. Membrane topology analysis of the sensor kinase KdpD of *Escherichia coli*. *J Biol Chem* 1995;270:28282–8.
- [158] Maier KS, Hubich S, Liebhart H, Krauss S, Kuhn A, Facey SJ. An amphiphilic region in the cytoplasmic domain of KdpD is recognized by the signal recognition particle and targeted to the *Escherichia coli* membrane. *Mol Microbiol* 2008;147:1–84.
- [159] Parltitz R, Eitan A, Stjepanovic G, Bahar I, Bange G, Bibi E, et al. *Escherichia coli* signal recognition particle receptor FtsY contains an essential and autonomous membrane-binding amphipathic helix. *J Biol Chem* 2007;282:32176–84.
- [160] Wimley WC, White SH. Membrane partitioning: distinguishing bilayer effects from the hydrophobic effect. *Biochemistry* 1993;32:6307–12.
- [161] Luecke H, Schobert B, Richter H-T, Cartailh J-P, Lanyi J. Structure of bacteriorhodopsin at 1.55 Å resolution. *J Mol Biol* 1999;291:899–911.
- [162] White SH, Ladokhin AS, Jayasinghe S, Hristova K. How membranes shape protein structure. *J Biol Chem* 2001;276:32395–8.
- [163] White SH, Wimley WC. Hydrophobic interactions of peptides with membrane interfaces. *Biochim Biophys Acta* 1998;1376:339–52.